

This Calculator format and content were approved
by the Ministry of Health in January 2020

Nplate[®] (romiplostim) Dose Calculator

**! Important: This calculator is ONLY to be used by
health care professionals.
This calculator must not be given to patients
who self-inject Nplate.**



Handling Guidance

- Nplate® can only be reconstituted with preservative-free sterile water for injections. **DO NOT** use sodium chloride solution for injection or bacteriostatic water for reconstitution of Nplate®.
- Nplate® is available:
 - 250 mcg on its own as a powder for solution for injections (paediatric and adult patients)
- Refrigerate (2° to 8°C) Nplate® and protect from light. Do not freeze.
- Nplate® should be used immediately after reconstitution. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 25°C or 24 hours in a refrigerator (2° to 8°C).
- Reconstituted Nplate® must also be protected from light.
- Any unused product or waste material should be disposed of in accordance with local requirements.
- Nplate® is a protein – **DO NOT SHAKE OR VIGOROUSLY AGITATE** the vial during reconstitution.

Steps for Reconstitution, Dilution (if Required) and Administration

Reconstitution:

Sterile water for injections only should be used when reconstituting the medicinal product. **DO NOT** use sodium chloride solutions for injection or bacteriostatic water for reconstitution of Nplate®.

1. Nplate® should be reconstituted in accordance with good aseptic practice.
2. **Sterile water for injections** should be injected into the vial.
3. The vial contents may be swirled gently and inverted during dissolution.
The vial should not be shaken or vigorously agitated. Generally, dissolution of Nplate® takes less than 2 minutes.
4. Visually inspect the solution for particulate matter and discolouration before administration.
 - The reconstituted solution should be clear and colourless.
 - The reconstituted solution should not be administered if particulate matter and/ or discolouration are observed.

Any unused product or waste material should be disposed of in accordance with local requirements.

Vial content:

Nplate® single-use vial	Total vial content of romiplostim		Volume of sterile water for injections		Deliverable product and volume	Final concentration
250 mcg	375 mcg	add	0.72 mL	=	250 mcg in 0.50 mL	500 mcg/mL

Steps for Reconstitution, Dilution (if Required) and Administration

Dilution (required when the calculated individual patient dose is less than 23 mcg):

Preservative-free, sterile, sodium chloride 9 mg/mL (0.9%) solution for injection only must be used for dilution.

DO NOT use dextrose (5%) in water or sterile water for injection for dilution.

Initial reconstitution of Nplate® with designated volumes of sterile water for injections results in a concentration of 500 mcg/mL in all vial sizes (see previous page and the vial content table). If the calculated individual patient dose is less than 23 mcg, an additional dilution step to 125 mcg/mL with **preservative-free, sterile, sodium chloride 9 mg/mL (0.9%) solution for injection** is required to ensure accurate dosing (see table below).

Dilution guidelines (refer to dose calculator for total injection volume to administer using the diluted concentration):

Nplate single-use vial	Add this volume of preservative-free, sterile, sodium chloride 9 mg/mL (0.9%) solution for injection to the reconstituted vial	Concentration after dilution
250 mcg	2.25 mL	125 mcg/mL

Storage of reconstituted and diluted Nplate®:

After reconstitution: chemical and physical in-use stability has been demonstrated for 24 hours at 25°C and for 24 hours at 2° to 8°C, when protected from light and **kept in the original vial**.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 25°C or 24 hours in a refrigerator (2° to 8°C), protected from light.

After dilution: chemical and physical in-use stability has been demonstrated for 4 hours at 25°C when the diluted product was held in a disposable syringe, or 4 hours in a refrigerator (2° to 8°C) when the diluted product was held in the original vial.

From a microbiological point of view, the diluted medicinal product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 4 hours at 25°C in disposable syringes, or 4 hours in a refrigerator (2° to 8°C) in the original vials, protected from light.


How to Calculate Nplate[®] Dose

Calculate initial dose:

1. Initial dose for Nplate[®] is 1 mcg/kg based on **actual body weight at initiation of therapy**.
2. Determine patient's weight in kilograms.
3. The individual patient dose does not have to be determined in order to use the dosing calculator.
4. Refer to dosing calculator windows on the following pages to achieve total injection volume required according to patient's weight (kg) and dose in mcg/kg.

Subsequent doses:

1. Determine patient's platelet count and previous week's dose.
2. Refer to dose adjustment table to determine how to adjust dose in mcg/kg based on changes in platelet counts.
3. In paediatric patients, future dose adjustments are based on changes in platelet counts **and** changes in body weight. Reassessment of body weight is recommended every 12 weeks.
4. Refer to dosing calculator windows on the following pages to achieve total injection volume required according to patient's weight (kg) and dose in mcg/kg.
5. Platelet counts should be assessed weekly until a stable platelet count ($\geq 50 \times 10^9/L$ for at least 4 weeks without dose adjustment) has been achieved. Platelet counts should be assessed monthly thereafter and appropriate dose adjustments made as per the dose adjustment table in order to maintain platelet counts within the recommended range.



Patients weighing 6 kg to 22 kg

Volumes to administer when no additional dilution is required

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Total injection volume (mL) required for individual patient dose										

Patients weighing 6 kg to 22 kg

Volumes to administer after additional dilution to 125 mcg/mL

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
Total injection volume (mL) required for individual patient dose										

When additional dilution is required, please refer to the window on the right on this page to achieve total injection volume required using the diluted product at 125 mcg/mL. Please note that calculation of individual patient dose is not required when using the dosing calculator window.

Individual patient dose (mcg) = patient's initial weight (kg) × dose in mcg/kg

Injection volume in mL* = $\frac{\text{individual patient dose (mcg)}}{500 \text{ mcg/mL}^{**}}$

*Round volume to the nearest hundredth mL

**Or 125 mcg/mL, if additional dilution required

 = Additional dilution required

Dilution guidelines:

Nplate® single-use vial	Add this volume of preservative-free, sterile, sodium chloride 9 mg/mL (0.9%) solution for injection to the reconstituted vial	Concentration after dilution
250 mcg	2.25 mL	125 mcg/mL

PULL

Patients weighing 6 kg to 22 kg

Volumes to administer when no additional dilution is required

6									
7									
8									
9									
10									
11									
12									
13									
14									
15									
16									
17									
18									
19									
20									
21									
22									

Patients weighing 6 kg to 22 kg

Volumes to administer after additional dilution to 125 mcg/mL

6	0.05	0.10	0.14
7	0.06	0.11	0.17
8	0.06	0.13	
9	0.07	0.14	
10	0.08	0.16	
11	0.09	0.18	
12	0.10		
13	0.10		
14	0.11		
15	0.12		
16	0.13		
17	0.14		
18	0.14		
19	0.15		
20	0.16		
21	0.17		
22	0.18		

PULL

Patients weighing 23kg to 45kg

Patients weighing 46kg to 68kg

23	0.05	0.09	0.14	0.18	0.23	0.28	0.32	0.37	0.41	0.46
24	0.05	0.10	0.14	0.19	0.24	0.29	0.34	0.38	0.43	0.48
25	0.05	0.10	0.15	0.20	0.25	0.30	0.35	0.40	0.45	0.50
26	0.05	0.10	0.16	0.21	0.26	0.31	0.36	0.42	0.47	0.52
27	0.05	0.11	0.16	0.22	0.27	0.32	0.38	0.43	0.49	0.54
28	0.06	0.11	0.17	0.22	0.28	0.34	0.39	0.45	0.50	0.56
29	0.06	0.12	0.17	0.23	0.29	0.35	0.41	0.46	0.52	0.58
30	0.06	0.12	0.18	0.24	0.30	0.36	0.42	0.48	0.54	0.60
31	0.06	0.12	0.19	0.25	0.31	0.37	0.43	0.50	0.56	0.62
32	0.06	0.13	0.19	0.26	0.32	0.38	0.45	0.51	0.58	0.64
33	0.07	0.13	0.20	0.26	0.33	0.40	0.46	0.53	0.59	0.66
34	0.07	0.14	0.20	0.27	0.34	0.41	0.48	0.54	0.61	0.68
35	0.07	0.14	0.21	0.28	0.35	0.42	0.49	0.56	0.63	0.70
36	0.07	0.14	0.22	0.29	0.36	0.43	0.50	0.58	0.65	0.72
37	0.07	0.15	0.22	0.30	0.37	0.44	0.52	0.59	0.67	0.74
38	0.08	0.15	0.23	0.30	0.38	0.46	0.53	0.61	0.68	0.76
39	0.08	0.16	0.23	0.31	0.39	0.47	0.55	0.62	0.70	0.78
40	0.08	0.16	0.24	0.32	0.40	0.48	0.56	0.64	0.72	0.80
41	0.08	0.16	0.25	0.33	0.41	0.49	0.57	0.66	0.74	0.82
42	0.08	0.17	0.25	0.34	0.42	0.50	0.59	0.67	0.76	0.84
43	0.09	0.17	0.26	0.34	0.43	0.52	0.60	0.69	0.77	0.86
44	0.09	0.18	0.26	0.35	0.44	0.53	0.62	0.70	0.79	0.88
45	0.09	0.18	0.27	0.36	0.45	0.54	0.63	0.72	0.81	0.90

46	0.09	0.18	0.28	0.37	0.46	0.55	0.64	0.74	0.83	0.92
47	0.09	0.19	0.28	0.38	0.47	0.56	0.66	0.75	0.85	0.94
48	0.10	0.19	0.29	0.38	0.48	0.58	0.67	0.77	0.86	0.96
49	0.10	0.20	0.29	0.39	0.49	0.59	0.69	0.78	0.88	0.98
50	0.10	0.20	0.30	0.40	0.50	0.60	0.70	0.80	0.90	1.00
51	0.10	0.20	0.31	0.41	0.51	0.61	0.71	0.82	0.92	1.02
52	0.10	0.21	0.31	0.42	0.52	0.62	0.73	0.83	0.94	1.04
53	0.11	0.21	0.32	0.42	0.53	0.64	0.74	0.85	0.95	1.06
54	0.11	0.22	0.32	0.43	0.54	0.65	0.76	0.86	0.97	1.08
55	0.11	0.22	0.33	0.44	0.55	0.66	0.77	0.88	0.99	1.10
56	0.11	0.22	0.34	0.45	0.56	0.67	0.78	0.90	1.01	1.12
57	0.11	0.23	0.34	0.46	0.57	0.68	0.80	0.91	1.03	1.14
58	0.12	0.23	0.35	0.46	0.58	0.70	0.81	0.93	1.04	1.16
59	0.12	0.24	0.35	0.47	0.59	0.71	0.83	0.94	1.06	1.18
60	0.12	0.24	0.36	0.48	0.60	0.72	0.84	0.96	1.08	1.20
61	0.12	0.24	0.37	0.49	0.61	0.73	0.85	0.98	1.10	1.22
62	0.12	0.25	0.37	0.50	0.62	0.74	0.87	0.99	1.12	1.24
63	0.13	0.25	0.38	0.50	0.63	0.76	0.88	1.01	1.13	1.26
64	0.13	0.26	0.38	0.51	0.64	0.77	0.90	1.02	1.15	1.28
65	0.13	0.26	0.39	0.52	0.65	0.78	0.91	1.04	1.17	1.30
66	0.13	0.26	0.40	0.53	0.66	0.79	0.92	1.06	1.19	1.32
67	0.13	0.27	0.40	0.54	0.67	0.80	0.94	1.07	1.21	1.34
68	0.14	0.27	0.41	0.54	0.68	0.82	0.95	1.09	1.22	1.36

FOLD

FOLD

How to Calculate Nplate® Dose

Volumes to administer for patients weighing 23 kg and above do not require additional dilution

Patients weighing 23kg to 45kg

Patients weighing 46kg to 68kg

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg
<input type="text"/>	<input type="text"/>									
Total injection volume (mL) required for individual patient dose										

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg
<input type="text"/>	<input type="text"/>									
Total injection volume (mL) required for individual patient dose										

Individual patient dose (mcg) = patient's initial weight (kg) × dose in mcg/kg

Injection volume in mL* = $\frac{\text{individual patient dose (mcg)}}{500 \text{ mcg/mL}}$

*Round volume to the nearest hundredth mL

FOLD and GLUE

FOLD

FOLD

How to Calculate Nplate® Dose

Volumes to administer for patients weighing 23 kg and above do not require additional dilution

Patients weighing 69kg to 91kg

Patients weighing 92 kg to 114 kg

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg
<input type="text"/>	<input type="text"/>									
Total injection volume (mL) required for individual patient dose										

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg
<input type="text"/>	<input type="text"/>									
Total injection volume (mL) required for individual patient dose										

Individual patient dose (mcg) = patient's initial weight (kg) × dose in mcg/kg

Injection volume in mL* = $\frac{\text{individual patient dose (mcg)}}{500 \text{ mcg/mL}}$

*Round volume to the nearest hundredth mL

FOLD and GLUE

PULL

Patients weighing 69kg to 91kg

Patients weighing 92kg to 114kg

69	0.14	0.28	0.41	0.55	0.69	0.83	0.97	1.10	1.24	1.38
70	0.14	0.28	0.42	0.56	0.70	0.84	0.98	1.12	1.26	1.40
71	0.14	0.28	0.43	0.57	0.71	0.85	0.99	1.14	1.28	1.42
72	0.14	0.29	0.43	0.58	0.72	0.86	1.01	1.15	1.30	1.44
73	0.15	0.29	0.44	0.58	0.73	0.88	1.02	1.17	1.31	1.46
74	0.15	0.30	0.44	0.59	0.74	0.89	1.04	1.18	1.33	1.48
75	0.15	0.30	0.45	0.60	0.75	0.90	1.05	1.20	1.35	1.50
76	0.15	0.30	0.46	0.61	0.76	0.91	1.06	1.22	1.37	1.52
77	0.15	0.31	0.46	0.62	0.77	0.92	1.08	1.23	1.39	1.54
78	0.16	0.31	0.47	0.62	0.78	0.94	1.09	1.25	1.40	1.56
79	0.16	0.32	0.47	0.63	0.79	0.95	1.11	1.26	1.42	1.58
80	0.16	0.32	0.48	0.64	0.80	0.96	1.12	1.28	1.44	1.60
81	0.16	0.32	0.49	0.65	0.81	0.97	1.13	1.30	1.46	1.62
82	0.16	0.33	0.49	0.66	0.82	0.98	1.15	1.31	1.48	1.64
83	0.17	0.33	0.50	0.66	0.83	1.00	1.16	1.33	1.49	1.66
84	0.17	0.34	0.50	0.67	0.84	1.01	1.18	1.34	1.51	1.68
85	0.17	0.34	0.51	0.68	0.85	1.02	1.19	1.36	1.53	1.70
86	0.17	0.34	0.52	0.69	0.86	1.03	1.20	1.38	1.55	1.72
87	0.17	0.35	0.52	0.70	0.87	1.04	1.22	1.39	1.57	1.74
88	0.18	0.35	0.53	0.70	0.88	1.06	1.23	1.41	1.58	1.76
89	0.18	0.36	0.53	0.71	0.89	1.07	1.25	1.42	1.60	1.78
90	0.18	0.36	0.54	0.72	0.90	1.08	1.26	1.44	1.62	1.80
91	0.18	0.36	0.55	0.73	0.91	1.09	1.27	1.46	1.64	1.82

92	0.18	0.37	0.55	0.74	0.92	1.10	1.29	1.47	1.66	1.84
93	0.19	0.37	0.56	0.74	0.93	1.12	1.30	1.49	1.67	1.86
94	0.19	0.38	0.56	0.75	0.94	1.13	1.32	1.50	1.69	1.88
95	0.19	0.38	0.57	0.76	0.95	1.14	1.33	1.52	1.71	1.90
96	0.19	0.38	0.58	0.77	0.96	1.15	1.34	1.54	1.73	1.92
97	0.19	0.39	0.58	0.78	0.97	1.16	1.36	1.55	1.75	1.94
98	0.20	0.39	0.59	0.78	0.98	1.18	1.37	1.57	1.76	1.96
99	0.20	0.40	0.59	0.79	0.99	1.19	1.39	1.58	1.78	1.98
100	0.20	0.40	0.60	0.80	1.00	1.20	1.40	1.60	1.80	2.00
101	0.20	0.40	0.61	0.81	1.01	1.21	1.41	1.62	1.82	2.02
102	0.20	0.41	0.61	0.82	1.02	1.22	1.43	1.63	1.84	2.04
103	0.21	0.41	0.62	0.82	1.03	1.24	1.44	1.65	1.85	2.06
104	0.21	0.42	0.62	0.83	1.04	1.25	1.46	1.66	1.87	2.08
105	0.21	0.42	0.63	0.84	1.05	1.26	1.47	1.68	1.89	2.10
106	0.21	0.42	0.64	0.85	1.06	1.27	1.48	1.70	1.91	2.12
107	0.21	0.43	0.64	0.86	1.07	1.28	1.50	1.71	1.93	2.14
108	0.22	0.43	0.65	0.86	1.08	1.30	1.51	1.73	1.94	2.16
109	0.22	0.44	0.65	0.87	1.09	1.31	1.53	1.74	1.96	2.18
110	0.22	0.44	0.66	0.88	1.10	1.32	1.54	1.76	1.98	2.20
111	0.22	0.44	0.67	0.89	1.11	1.33	1.55	1.78	2.00	2.22
112	0.22	0.45	0.67	0.90	1.12	1.34	1.57	1.79	2.02	2.24
113	0.23	0.45	0.68	0.90	1.13	1.36	1.58	1.81	2.03	2.26
114	0.23	0.46	0.68	0.91	1.14	1.37	1.60	1.82	2.05	2.28

PULL

Patients weighing 115kg to 137 kg

115	0.23	0.46	0.69	0.92	1.15	1.38	1.61	1.84	2.07	2.30
116	0.23	0.46	0.70	0.93	1.16	1.39	1.62	1.86	2.09	2.32
117	0.23	0.47	0.70	0.94	1.17	1.40	1.64	1.87	2.11	2.34
118	0.24	0.47	0.71	0.94	1.18	1.42	1.65	1.89	2.12	2.36
119	0.24	0.48	0.71	0.95	1.19	1.43	1.67	1.90	2.14	2.38
120	0.24	0.48	0.72	0.96	1.20	1.44	1.68	1.92	2.16	2.40
121	0.24	0.48	0.73	0.97	1.21	1.45	1.69	1.94	2.18	2.42
122	0.24	0.49	0.73	0.98	1.22	1.46	1.71	1.95	2.20	2.44
123	0.25	0.49	0.74	0.98	1.23	1.48	1.72	1.97	2.21	2.46
124	0.25	0.50	0.74	0.99	1.24	1.49	1.74	1.98	2.23	2.48
125	0.25	0.50	0.75	1.00	1.25	1.50	1.75	2.00	2.25	2.50
126	0.25	0.50	0.76	1.01	1.26	1.51	1.76	2.02	2.27	2.52
127	0.25	0.51	0.76	1.02	1.27	1.52	1.78	2.03	2.29	2.54
128	0.26	0.51	0.77	1.02	1.28	1.54	1.79	2.05	2.30	2.56
129	0.26	0.52	0.77	1.03	1.29	1.55	1.81	2.06	2.32	2.58
130	0.26	0.52	0.78	1.04	1.30	1.56	1.82	2.08	2.34	2.60
131	0.26	0.52	0.79	1.05	1.31	1.57	1.83	2.10	2.36	2.62
132	0.26	0.53	0.79	1.06	1.32	1.58	1.85	2.11	2.38	2.64
133	0.27	0.53	0.80	1.06	1.33	1.60	1.86	2.13	2.39	2.66
134	0.27	0.54	0.80	1.07	1.34	1.61	1.88	2.14	2.41	2.68
135	0.27	0.54	0.81	1.08	1.35	1.62	1.89	2.16	2.43	2.70
136	0.27	0.54	0.82	1.09	1.36	1.63	1.90	2.18	2.45	2.72
137	0.27	0.55	0.82	1.10	1.37	1.64	1.92	2.19	2.47	2.74

Patients weighing 138 kg to 150 kg

138	0.28	0.55	0.83	1.10	1.38	1.66	1.93	2.21	2.48	2.76
139	0.28	0.56	0.83	1.11	1.39	1.67	1.95	2.22	2.50	2.78
140	0.28	0.56	0.84	1.12	1.40	1.68	1.96	2.24	2.52	2.80
141	0.28	0.56	0.85	1.13	1.41	1.69	1.97	2.26	2.54	2.82
142	0.28	0.57	0.85	1.14	1.42	1.70	1.99	2.27	2.56	2.84
143	0.29	0.57	0.86	1.14	1.43	1.72	2.00	2.29	2.57	2.86
144	0.29	0.58	0.86	1.15	1.44	1.73	2.02	2.30	2.59	2.88
145	0.29	0.58	0.87	1.16	1.45	1.74	2.03	2.32	2.61	2.90
146	0.29	0.58	0.88	1.17	1.46	1.75	2.04	2.34	2.63	2.92
147	0.29	0.59	0.88	1.18	1.47	1.76	2.06	2.35	2.65	2.94
148	0.30	0.59	0.89	1.18	1.48	1.78	2.07	2.37	2.66	2.96
149	0.30	0.60	0.89	1.19	1.49	1.79	2.09	2.38	2.68	2.98
150	0.30	0.60	0.90	1.20	1.50	1.80	2.10	2.40	2.70	3.00

How to Calculate Nplate[®] Dose

Volumes to administer for patients weighing 23 kg and above do not require additional dilution

Patients weighing 115 kg to 137 kg

Patients weighing 138 kg to 150 kg

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg
<input type="text"/>	<input type="text"/>									
Total injection volume (mL) required for individual patient dose										

Patient weight (kg)	Dose (mcg/kg)									
	1	2	3	4	5	6	7	8	9	10
	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg	mcg/kg
<input type="text"/>	<input type="text"/>									
Total injection volume (mL) required for individual patient dose										

Individual patient dose (mcg) = patient's initial weight (kg) × dose in mcg/kg

Injection volume in mL* = $\frac{\text{individual patient dose (mcg)}}{500 \text{ mcg/mL}}$

*Round volume to the nearest hundredth mL

Dosing Adjustments

Dose adjustment guidance based on platelet counts:

Platelet count ($\times 10^9/L$)	Action
< 50	Increase once weekly dose by 1 mcg/kg
> 150 for two consecutive weeks	Decrease once weekly dose by 1 mcg/kg
> 250	Do not administer, continue to assess the platelet count weekly After the platelet count has fallen to $< 150 \times 10^9/L$, resume dosing with once weekly dose reduced by 1 mcg/kg

Due to the interindividual variable platelet response, in some patients platelet count may abruptly fall below $50 \times 10^9/L$ after dose reduction or treatment discontinuation. In these cases, if clinically appropriate, higher cut-off levels of platelet count for dose reduction ($200 \times 10^9/L$) and treatment interruption ($400 \times 10^9/L$) may be considered according to medical judgement.

A maximum once weekly dose of 10 mcg/kg should not be exceeded.

Nplate[®] should continue to be administered weekly unless a platelet count of $> 250 \times 10^9/L$ is achieved.

Directions for Dose Reconstitution and Administration

- Nplate® is a highly potent peptide administered subcutaneously as a low-volume dose. Nplate® is supplied as a powder for reconstitution with sterile water for injections.
- When reconstituted as directed, the vial provides a solution with a concentration of 500 mcg/mL . Because of the hold up in the vial and syringe, the deliverable amount of Nplate® is less than the reconstituted amount.
 - Nplate® 250 micrograms powder for solution for injection should be reconstituted with 0.72 mL sterile water for injections, yielding a deliverable volume of 0.50 mL . An additional overfill is included in each vial to ensure that 250 mcg of romiplostim can be delivered (see vial content table below).

Vial content:

Nplate® single-use vial	Total vial content of romiplostim		Volume of sterile water for injections		Deliverable product and volume	Final concentration
250 mcg	375 mcg	add	0.72 mL	=	250 mcg in 0.50 mL	500 mcg/mL

- Smaller amounts of Nplate® may be required for weekly injection in patients for whom low weekly doses are appropriate, as well as during the titration phase.
- Administer Nplate® as a weekly subcutaneous injection with dose adjustments based on the platelet count response (paediatric and adult patients) and patient's weight (paediatric patients only).
- Injection volume may be very small. Use a syringe with graduations to 0.01 mL.

Prescribing Information

Reporting suspect adverse reactions after authorisation of the medicinal product is important.
It allows continued monitoring of the benefit/risk balance of the medicinal product.

Healthcare professionals are asked to report any suspected adverse reactions.

Adverse events can be reported to the Ministry of Health via
<https://sideeffects.health.gov.il/>

Or reported to local distributor Medison: pv@Medison.co.il

If you have any questions about the administration of NPLATE refer to the Israeli PI







NPL-35-01-0220