

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Glypressin

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each ampoule contains terlipressin acetate 1 mg.

Excipient: Each 5 ml vial of solvent contains 17.7 mg sodium.

For full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder and solvent for solution for injection.

White, sterile powder.

Clear, colourless solvent.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Bleeding oesophageal varices.

Treatment of type I hepatorenal syndrome.

4.2 Posology and method of administration

1. Oesophageal varices bleeding

Unless otherwise prescribed, initially IV injection of 1-2mg terlipressin acetate, equivalent to 1-2 vials of Glypressin 1mg, is slowly administered to adults. The maintenance dose is 1 mg terlipressin acetate , equivalent to 1 vial of Glypressin 1mg after 4-6 hours. The standard value of the maximum daily dose of Glypressin is 120-150 ug/kg body weight. For an adult person of 70 kg body weight, this corresponds to a dose of 8- 10 vials per day, to be administered in 4-hour intervals.

2) In type 1 hepatorenal syndrome:

3 to 4 mg every 24 hours as 3 or 4 administrations.

In the absence of any reduction of the serum creatinine after 3 days of treatment, cessation of Glypressin treatment is advised.

In the other cases, Glypressin treatment is to be pursued until the obtaining either of a serum creatinine less than 130 µmol/litre or of a drop of at least 30 % in the serum creatinine with respect to the value measured at the time of diagnosis of hepatorenal syndrome. The standard average duration of treatment is 10 days.

Special Populations

Elderly patients

There is no data available regarding dosage recommendation in the elderly.

Paediatric population

There is no data available regarding dosage recommendation in the paediatric population.

Method of Administration

IV injection

4.3 Contraindications

Hypersensitivity to terlipressin or any other excipient of the product.

Contraindicated in pregnancy.

4.4 Special warnings and precautions for use

Cardiac, pulmonary and vascular disease

During treatment regular monitoring and control of blood pressure, ECG, heart rate, serum levels of sodium and potassium, as well as fluid balance are required.

Caution should be exercised in treating patients with hypertension, recognised heart disease, renal dysfunction, cerebral or peripheral vascular disease, asthma or respiratory failure.

Injection site reaction To avoid local necrosis at the injection site, the injection must be administered intravenously.

Septic shock

In patients with septic shock with a low cardiac output terlipressin should not be used.

Torsade de pointes

During clinical trials and post-marketing experience, several cases of QT interval prolongation and ventricular arrhythmias including “Torsade de pointes” have been reported (see section 4.8). In most cases, patients had predisposing factors such as basal prolongation of the QT interval, electrolyte abnormalities (hypokalemia, hypomagnesemia) or medications with concomitant effect on QT prolongation. Therefore, extreme caution should be exercised in the use of terlipressin in patients with a history of QT interval prolongation, electrolyte abnormalities, or concomitant medications that can prolong the QT interval (see section 4.5).

Prior to use of terlipressin for hepatorenal syndrome, it must be ascertained that the patient has an acute functional renal failure and this functional renal failure does not respond to a suitable plasma expansion therapy.

Paediatric population and elderly patients: Particular caution should be exercised in the treatment of children and elderly patients, as experience is limited in these groups.

There is no data available regarding dosage recommendation in these special patient categories.

Excipients

This medicine contains less than 1 mmol sodium (23 mg) per vial, that is to say essentially ‘sodium-free’.

4.5 Interaction with other medicinal products and other forms of interaction

The hypotensive effect of non-selective beta-blockers on the portal vein is increased with terlipressin.

Concomitant treatment with medicinal products with a known bradycardiac effect (e.g. propofol, sufentanil) may lower the heart rate and cardiac output. These effects are due to reflexogenic inhibition of the cardiac activity via the vagus nerve due to the elevated blood pressure.

Terlipressin can trigger “torsade de pointes” (see sections 4.4 and 4.8). Therefore, extreme caution should be exercised in the use of terlipressin in patients with concomitant medications that can prolong the QT interval, such as class IA and III antiarrhythmics, erythromycin, certain antihistamines and tricyclic antidepressants or medications that may cause hypokalaemia or hypomagnesemia (e.g. some diuretics).

4.6 Pregnancy and lactation

Pregnancy

Treatment with GLYPRESSIN during pregnancy is contraindicated (see sections 4.3 and 5.3).

GLYPRESSIN has been shown to cause uterine contractions and increased intrauterine pressure in early pregnancy and may decrease uterine blood flow. GLYPRESSIN may have harmful effects on pregnancy and foetus.

Spontaneous abortion and malformation have been shown in rabbits after treatment with GLYPRESSIN.

Breast-feeding

It is not known whether GLYPRESSIN is excreted in human breast milk. The excretion of GLYPRESSIN in milk has not been studied in animals. A risk to the suckling child cannot be excluded. A decision on whether to continue/discontinue breast-feeding or to continue/discontinue therapy with GLYPRESSIN should be made taking into account the benefit of breast-feeding to the child and the benefit of GLYPRESSIN therapy to the woman.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

4.8 Undesirable effects

The most commonly reported undesired effects in clinical trials are paleness, increased blood pressure, abdominal pain, nausea, diarrhoea, and headache.

Tabulated list of adverse reactions\

MedDRA System Organ Class	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)
Metabolism and nutrition disorders		Hyponatraemia	
Nervous system disorders	Headache		
Cardiac disorders	Bradycardia	Atrial fibrillation Ventricular extracystoles Tachycardia Myocardial infarction Torsade de pointes Cardiac failure cyanosis	
Vascular disorders	vasoconstriction Peripheral ischemia pallor Hypertension	Hot flush	
Respiratory, thoracic and mediastinal disorders		Respiratory distress Pulmonary oedema Respiratory failure	Dyspnoea
Gastrointestinal disorders	abdominal pain diarrhoea	nausea Intestinal ischemia vomiting	
Skin and subcutaneous tissue disorders		Skin necrosis	
Pregnancy, puerperium and perinatal conditions		Uterine hypertonus Uterine ischemia	
General disorders and administration site disorders		Injection site necrosis Chest pain	

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il/>

4.9 Overdose

The recommended dose in the specific patient population should not be exceeded as the risk of severe circulatory adverse effects is dose-dependent.

Elevated blood pressure in patients with hypertension can be controlled with clonidine 150 µg IV. Severe bradycardia should be treated with atropine

5. PHARMACOLOGICAL PROPERTIES

5.0 Pharmacotherapeutic group

Pharmacotherapeutic group: Posterior pituitary lobe hormones (vasopressin and analogues)

ATC code: H01B A04

Terlipressin (Triglycyl-Lysine-Vasopressin) is a synthetic analogue of the natural posterior pituitary hormone vasopressin.

Terlipressin is a pro-drug with partial, intrinsic activity by itself. Terlipressin is transformed into the fully active metabolite lysine-vasopressin (LVP) by enzymatic cleavage. LVP remains within the therapeutic concentration range over a period of 4-6 hours.

Doses of 1 and 2 mg terlipressin acetate effectively reduce the portal venous pressure and produce marked vasoconstriction. The lowering of portal pressure and azygos blood flow is dependent on dose. The effect of the low dose is reduced after 3 hours, while haemodynamic data show that 2 mg terlipressin acetate is more effective than 1 mg terlipressin acetate with a sustained effect throughout the treatment period (4 to 6 hours).

5.1 Pharmacodynamic properties

Glypressin may be regarded as a circulating depot of lysine vasopressin. Following i.v. injection, three glycyl moieties are enzymatically cleaved from the N terminal to release lysine vasopressin.

The slowly released vasopressin reduces blood flow in the splanchnic circulation in a prolonged manner, thereby helping to control bleeding from ruptured oesophageal varices.

5.2 Pharmacokinetic properties

Glypressin is administered by bolus IV injection. It shows a biphasic plasma level curve which indicates that a two-compartment model can be applied. The half-life of elimination ($T_{1/2 \beta}$) is about 50-70 minutes.

Lysine vasopressin reaches maximum plasma levels about 1-2 hours following i.v. administration and has a duration of activity of 4-6 hours.

5.3 Preclinical safety data

There are no pre-clinical data of relevance to the prescriber which are additional to that already included in other sections of the SPC.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Powder for solution for injection (vial):

Mannitol Hydrochloric acid (for pH adjustment)

Solvent (ampoule):

Sodium chloride

Hydrochloric acid (for pH adjustment)

Water for injections

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials

6.4 Special precautions for storage

Do not store above 25°C. Store in the original package in order to protect from light.

The reconstituted solution must be used immediately after reconstitution.

6.5 Nature and contents of container

Type I glass vial with bromobutyl rubber stopper and type I glass ampoule.

The vial contains terlipressin acetate 1 mg. The ampoule has a coloured dot indicating the cut area and contains 5 ml of clear, colourless solvent.

Pack size. The package contains 5 sets of 1 vial + 1 ampoule

6.6 Special precautions for disposal and other handling

Reconstitute Glypressin with the solvent provided immediately prior to use. For single use only. Discard any unused solution.

7. Manufacturer:

Ferring GmbH, Germany.

8. License Holder:

Ferring Pharmaceuticals Ltd., 8 Hashita St., P.O.Box 3551, Caesarea Industrial Park, 3088900

10. Registration number: 133- 74-31265

Revised in July 2020