

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Fampyra 10 mg prolonged-release tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each prolonged-release tablet contains 10 mg of fampridine.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Prolonged-release tablet.

off-white, film coated, oval biconvex 13 x 8 mm tablet with flat edge debossed with A10 on one side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Fampyra is indicated for the improvement of walking in adult patients with multiple sclerosis with walking disability (EDSS 4-7).

4.2 Posology and method of administration

Treatment with Fampyra is restricted to prescription and supervision by physicians experienced in the management of MS.

Posology

The recommended dose is one 10 mg tablet, twice daily, taken 12 hours apart (one tablet in the morning and one tablet in the evening). Fampyra should not be administered more frequently or at higher doses than recommended (see section 4.4). The tablets should be taken without food (see section 5.2).

Starting and Evaluating Fampyra Treatment

- Initial prescription should be limited to two to four weeks of therapy as clinical benefits should generally be identified within two to four weeks after starting Fampyra
- An assessment of walking ability, e.g. the Timed 25 Foot Walk (T25FW) or Twelve Item Multiple Sclerosis Walking Scale (MSWS-12), is recommended to evaluate improvement within two to four weeks. If no improvement is observed, Fampyra should be discontinued
- Fampyra should be discontinued if benefit is not reported by patients.

Re-Evaluating Fampyra Treatment

If decline in walking ability is observed, physicians should consider an interruption to treatment in order to reassess the benefits of Fampyra (see above). The re-evaluation should include withdrawal of Fampyra and performing an assessment of walking ability. Fampyra should be discontinued if

patients no longer receive walking benefit.

Missed Dose

The usual dosing regimen should always be followed. A double dose should not be taken if a dose is missed.

Older people

Renal function should be checked in older people before starting treatment with Fampyra. Monitoring renal function to detect any renal impairment is recommended in older people (see section 4.4).

Patients with renal impairment

Fampyra is contraindicated in patients with mild, moderate and severe renal impairment (creatinine clearances <80 ml/min) (see section 4.3).

Patients with hepatic impairment

No dose adjustment is required for patients with hepatic impairment.

Paediatric population

Fampyra is not indicated for children and adolescents under 18 years old
The safety and efficacy of Fampyra in children aged 0 to 18 years have not been established. No data are available.

Method of Administration

Fampyra is for oral use.

The tablet must be swallowed whole. It must not be divided, crushed, dissolved, sucked or chewed.

4.3 Contraindications

Hypersensitivity to fampridine or to any of the excipients listed in section 6.1.

Concurrent treatment with other medicinal products containing fampridine (4-aminopyridine).

Patients with prior history or current presentation of seizure.

Patients with mild, moderate or severe renal impairment (creatinine clearances <80 ml/min).

Concomitant use of Fampyra with medicinal products that are inhibitors of Organic Cation Transporter 2 (OCT2) for example, cimetidine.

4.4 Special warnings and precautions for use

Seizure risk

Treatment with fampridine increases seizure risk (see section 4.8).

Fampyra should be administered with caution in the presence of any factors which may lower seizure threshold.

Fampyra should be discontinued in patients who experience a seizure while on treatment.

Renal impairment

Fampyra is primarily excreted unchanged by the kidneys. Patients with renal impairment have higher plasma concentrations which are associated with increased adverse reactions, in particular

neurological effects. Determining renal function before treatment and its regular monitoring during treatment is recommended in all patients (particularly in older people in whom renal function might be reduced). Creatinine clearance can be estimated using the Cockcroft-Gault formula.

Fampyra should not be administered to patients with renal impairment (creatinine clearance <80 ml/min) (see section 4.3).

Caution is required when Fampyra is prescribed concurrently with medicinal products that are substrates of OCT2 for example, carvedilol, propranolol and metformin.

Hypersensitivity Reactions

In post-marketing experience, serious hypersensitivity reactions (including anaphylactic reaction) have been reported, the majority of these cases occurred within the first week of treatment. Particular attention should be given to patients with a previous history of allergic reactions. If an anaphylactic or other serious allergic reaction occurs, Fampyra should be discontinued and not restarted.

Other warnings and precautions

Fampyra should be administered with caution to patients with cardiovascular symptoms of rhythm and sinoatrial or atrioventricular conduction cardiac disorders (these effects are seen in overdose). There is limited safety information in these patients.

The increased incidence of dizziness and balance disorder seen with Fampyra may result in an increased risk of falls. Therefore, patients should use walking aids as needed.

In clinical studies low white blood cell counts were seen in 2.1% of Fampyra patients versus 1.9% of patients on placebo. Infections were seen in the clinical studies (see section 4.8) and increased infection rate and impairment of the immune response cannot be excluded.

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

Concurrent treatment with other medicinal products containing fampridine (4-aminopyridine) is contraindicated (see section 4.3).

Fampridine is eliminated mainly via the kidneys with active renal secretion accounting for about 60% (see section 5.2). OCT2 is the transporter responsible for the active secretion of fampridine. Thus, the concomitant use of fampridine with medicinal products that are inhibitors of OCT2 for example, cimetidine are contraindicated (see section 4.3) and concomitant use of fampridine with medicinal products that are substrates of OCT2 for example, carvedilol, propranolol and metformin is cautioned (see section 4.4.)

Interferon: fampridine has been administered concomitantly with interferon-beta and no pharmacokinetic medicinal product interactions were observed.

Baclofen: fampridine has been administered concomitantly with baclofen and no pharmacokinetic medicinal product interactions were observed.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are limited amount of data from the use of fampridine in pregnant women.

Animal studies have shown reproductive toxicity (see section 5.3). As a precautionary measure it is

preferable to avoid the use of Fampyra in pregnancy.

Breast-feeding

It is unknown whether fampridine is excreted in human or animal milk. Fampyra is not recommended during breast-feeding.

Fertility

In animal studies no effects on fertility were seen.

4.7 Effects on ability to drive and use machines

Fampyra has a moderate influence on the ability to drive and use machines because Fampyra can cause dizziness.

4.8 Undesirable effects

The safety of Fampyra has been evaluated in randomised controlled clinical studies, in open label long term studies and in the post marketing setting.

Adverse reactions identified are mostly neurological and include seizure, insomnia, anxiety, balance disorder, dizziness, paraesthesia, tremor, headache and asthenia. This is consistent with fampridine's pharmacological activity. The highest incidence of adverse reactions identified from placebo-controlled trials in multiple sclerosis patients with Fampyra given at the recommended dose, are reported as urinary tract infection (in approximately 12% of patients).

Adverse reactions are presented below by system organ class and absolute frequency. Frequencies are defined as: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

MedDRA SOC	Adverse Reaction	Frequency category
Infections and infestations	Urinary tract infection ¹	Very Common
	Influenza ¹	Common
	Nasopharyngitis ¹	Common
	Viral infection ¹	Common
Immune system disorders	Anaphylaxis	Uncommon
	Angioedema	Uncommon
	Hypersensitivity	Uncommon
Psychiatric disorders	Insomnia	Common
	Anxiety	Common
Nervous system disorders	Dizziness	Common
	Headache	Common
	Balance disorder	Common
	Paraesthesia	Common
	Tremor	Common
	Seizure	Uncommon
	Exacerbation of trigeminal neuralgia	Uncommon
Cardiac disorders	Palpitations	Common
	Tachycardia	Uncommon
Vascular disorders	Hypotension ²	Uncommon
Respiratory, thoracic and mediastinal disorders	Dyspnoea	Common
	Pharyngolaryngeal pain	Common
Gastrointestinal disorders	Nausea	Common
	Vomiting	Common
	Constipation	Common
	Dyspepsia	Common
Skin and subcutaneous tissue disorders	Rash	Uncommon
	Urticaria	Uncommon
Musculoskeletal and connective tissue disorders	Back pain	Common
General disorders and administration site conditions	Asthenia	Common
	Chest discomfort ²	Uncommon

¹ See section 4.4

² These symptoms were observed in the context of hypersensitivity

Description of selected adverse reactions

Seizure

In post-marketing experience, there have been reports of seizure, the frequency is not known (cannot be estimated from the available data). For further information on seizure risk, please refer to sections 4.3 and 4.4.

Hypersensitivity

In post-marketing experience, there have been reports of hypersensitivity reactions (including anaphylaxis) which have occurred with one or more of the following: dyspnoea, chest discomfort, hypotension, angioedema, rash and urticaria. For further information on hypersensitivity reactions, please refer to sections 4.3 and 4.4.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form <https://sideeffects.health.gov.il/>

4.9 Overdose

Symptoms

Acute symptoms of overdose with Fampyra were consistent with central nervous system excitation and included confusion, tremulousness, diaphoresis, seizure, and amnesia.

Central nervous system side effects at high doses of 4-aminopyridine include confusion, seizures, status epilepticus, involuntary and choreoathetoid movements. Other side effects at high doses include cases of cardiac arrhythmias (for example, supraventricular tachycardia and bradycardia) and ventricular tachycardia as a consequence of potential QT prolongation. Reports of hypertension have also been received.

Management

Patients who overdose should be provided supportive care. Repeated seizure activity should be treated with benzodiazepine, phenytoin, or other appropriate acute anti-seizure therapy.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other nervous system drugs, ATC code: N07XX07.

Pharmacodynamic effects

Fampyra is a potassium channel blocker. By blocking potassium channels, Fampyra reduces the leakage of ionic current through these channels, thereby prolonging repolarization and thus enhancing action potential formation in demyelinated axons and neurological function. Presumably, by enhancing action potential formation, more impulses might be conducted in the central nervous system.

Clinical efficacy and safety

Three phase III, randomised, double-blind, placebo controlled confirmatory studies, (MS-F203 and MS-F204 and 218MS305) have been performed. The proportion of responders was independent of concomitant immunomodulatory therapy (including interferons, glatiramer acetate, fingolimod and natalizumab). The Fampyra dose was 10 mg BID.

Studies MS-F203 and MS-F204

The primary endpoint in studies MS-F203 and MS-F204 was the responder rate in walking speed as measured by the Timed 25-foot Walk (T25FW). A responder was defined as a patient who consistently had a faster walking speed for at least three visits out of a possible four during the double blind period as compared to the maximum value among five off-treatment visits.

A significantly greater proportion of Fampyra treated patients were responders as compared to placebo (MS-F203: 34.8% vs. 8.3%, $p < 0.001$; MS-F204: 42.9% vs. 9.3%, $p < 0.001$).

Patients who responded to Fampyra increased their walking speed on average by 26.3% vs 5.3% on placebo ($p < 0.001$) (MS-F203) and 25.3% vs 7.8% ($p < 0.001$) (MS-F204). The improvement appeared

rapidly (within weeks) after starting Fampyra.

Statistically and clinically meaningful improvements in walking were seen, as measured by the 12-item Multiple Sclerosis Walking Scale.

1

Table 1: Studies MS-F203 and MS-F204

STUDY *	MS-F203		MS-F204	
	Placebo	Fampyra 10 mg BID	Placebo	Fampyra 10 mg BID
n of subjects	72	224	118	119
Consistent improvement	8.3%	34.8%	9.3%	42.9%
Difference		26.5%		33.5%
CI _{95%}		17.6%, 35.4%		23.2%, 43.9%
P-value		< 0.001		< 0.001
≥20% improvement	11.1%	31.7%	15.3%	34.5%
Difference		20.6%		19.2%
CI _{95%}		11.1%, 30.1%		8.5%, 29.9%
P-value		<0.001		<0.001
Walking speed Feet/sec	Ft per sec	Ft per sec	Ft per sec	Ft per sec
Baseline	2.04	2.02	2.21	2.12
Endpoint	2.15	2.32	2.39	2.43
Change	0.11	0.30	0.18	0.31
Difference		0.19		0.12
p-value		0.010		0.038
Average % Change	5.24	13.88	7.74	14.36
Difference		8.65		6.62
p-value		< 0.001		0.007
MSWS-12-score (mean, sem)				
Baseline	69.27 (2.22)	71.06 (1.34)	67.03 (1.90)	73.81 (1.87)
Average change	-0.01 (1.46)	-2.84 (0.878)	0.87 (1.22)	-2.77 (1.20)
Difference		2.83		3.65
p-value		0.084		0.021
LEMMT (mean, sem) (Lower Extremity Manual Muscle Test)				
Baseline	3.92 (0.070)	4.01 (0.042)	4.01 (0.054)	3.95 (0.053)
Average change	0.05 (0.024)	0.13 (0.014)	0.05 (0.024)	0.10 (0.024)
Difference		0.08		0.05
p-value		0.003		0.106
Ashworth Score (A test for muscle spasticity)				
Baseline	0.98 (0.078)	0.95 (0.047)	0.79 (0.058)	0.87 (0.057)
Average change	-0.09 (0.037)	-0.18 (0.022)	-0.07 (0.033)	-0.17 (0.032)
Difference		0.10		0.10
p-value		0.021		0.015

Study 218MS305

Study 218MS305 was conducted in 636 subjects with multiple sclerosis and walking disability. Duration of double-blind treatment was 24 weeks with a 2 week post-treatment follow-up. The primary endpoint was improvement in walking ability, measured as the proportion of patients achieving a mean improvement of ≥ 8 points from baseline MSWS-12 score over 24 weeks. In this study there was a statistically significant treatment difference, with a greater proportion of Fampyra treated patients demonstrating an improvement in walking ability, compared to placebo-controlled patients (relative risk of 1.38 (95% CI: [1.06, 1.70])). Improvements generally appeared within 2 to 4 weeks of initiation of treatment, and disappeared within 2 weeks of treatment cessation.

Fampyra treated patients also demonstrated a statistically significant improvement in the Timed Up and Go (TUG) test, a measure of static and dynamic balance and physical mobility. In this secondary endpoint, a greater proportion of Fampyra treated patients achieved $\geq 15\%$ mean improvement from baseline TUG speed over a 24 week period, compared to placebo. The difference in the Berg Balance Scale (BBS; a measure of static balance), was not statistically significant.

In addition, patients treated with Fampyra demonstrated a statistically significant mean improvement from baseline compared to placebo in the Multiple Sclerosis Impact Scale (MSIS-29) physical score (LSM difference -3.31, $p < 0.001$).

Table 2: Study 218MS305

Over 24 weeks	Placebo N = 318*	Fampyra 10 mg BID N = 315*	Difference (95% CI) <i>p</i> - value
Proportion of patients with mean improvement of ≥ 8 points from baseline MSWS-12 score	34%	43%	Risk difference: 10.4% (3% ; 17.8%) 0.006
MSWS-12 score Baseline Improvement from baseline	65.4 -2.59	63.6 -6.73	LSM: -4.14 (-6.22 ; -2.06) <0.001
TUG Proportion of patients with mean improvement of $\geq 15\%$ in TUG speed	35%	43%	Risk difference: 9.2% (0.9% ; 17.5%) 0.03
TUG Baseline Improvement from baseline (sec)	27.1 -1.94	24.9 -3.3	LSM: -1.36 (-2.85 ; 0.12) 0.07
MSIS-29 physical score Baseline Improvement from baseline	55.3 -4.68	52.4 -8.00	LSM: -3.31 (-5.13 ; -1.50) <0.001
BBS score Baseline Improvement from baseline	40.2 1.34	40.6 1.75	LSM: 0.41 (-0.13 ; 0.95) 0.141

*Intent to treat population = 633; LSM = Least square mean

5.2 Pharmacokinetic properties

Absorption:

Orally administered fampridine is rapidly and completely absorbed from the gastrointestinal tract. Fampridine has a narrow therapeutic index. Absolute bioavailability of Fampyra prolonged-release tablets has not been assessed, but relative bioavailability (as compared to an aqueous oral solution) is 95%. The Fampyra prolonged-release tablet has a delay in the absorption of fampridine manifested by slower rise to a lower peak concentration, without any effect on the extent of absorption.

When Fampyra tablets are taken with food, the reduction in the area under the plasma concentration-time curve ($AUC_{0-\infty}$) of fampridine is approximately 2-7% (10 mg dose). The small reduction in AUC is not expected to cause a reduction in the therapeutic efficacy. However, C_{max} increases by 15-23%. Since there is a clear relationship between C_{max} and dose related adverse reactions, it is recommended to take Fampyra without food (see section 4.2).

Distribution:

Fampridine is a lipid-soluble medicinal product which readily crosses the blood-brain barrier. Fampridine is largely unbound to plasma proteins (bound fraction varied between 3-7% in human plasma). Fampridine has a volume of distribution of approximately 2.6 l/kg.

Fampridine is not a substrate for P-glycoprotein.

Biotransformation:

Fampridine is metabolised in humans by oxidation to 3-hydroxy-4-aminopyridine and further conjugated to the 3-hydroxy-4-aminopyridine sulfate. No pharmacological activity was found for the fampridine metabolites against selected potassium channels *in vitro*.

The 3-hydroxylation of fampridine to 3-hydroxy-4-aminopyridine by human liver microsomes appeared to be catalyzed by Cytochrome P450 2E1 (CYP2E1).

There was evidence of direct inhibition of CYP2E1 by fampridine at 30 μ M (approximately 12% inhibition) which is approximately 100 times the average plasma fampridine concentration measured for the 10 mg tablet.

Treatment of cultured human hepatocytes with fampridine had little or no effect on induction of CYP1A2, CYP2B6, CYP2C9, CYP2C19, CYP2E1 or CYP3A4/5 enzyme activities.

Elimination:

The major route of elimination for fampridine is renal excretion, with approximately 90% of the dose recovered in urine as parent medicinal product within 24 hours. Renal clearance (CLR 370 ml/min) is substantially greater than glomerular filtration rate due to combined glomerular filtration and active excretion by the renal OCT2 transporter. Faecal excretion accounts for less than 1% of the administered dose.

Fampyra is characterized by linear (dose-proportional) pharmacokinetics with a terminal elimination half-life of approximately 6 hours. The maximum plasma concentration (C_{max}) and, to a smaller extent, area under the plasma concentration-time curve (AUC) increase proportionately with dose.

There is no evidence of clinically relevant accumulation of fampridine taken at the recommended dose in patients with full renal function. In patients with renal impairment, accumulation occurs relative to the degree of impairment.

Special Populations

Older people:

Clinical studies of Fampyra did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger patients. Fampyra is primarily excreted unchanged by the kidneys, and with creatinine clearance known to decrease with age, monitoring of renal function in older patients should be considered (see section 4.2).

Paediatric Population:

No data are available.

Patients with renal impairment:

Fampridine is eliminated primarily by the kidneys as unchanged medicinal product and therefore renal function should be checked in patients where renal function might be compromised. Patients with mild renal impairment can be expected to have approximately 1.7 to 1.9 times the fampridine concentrations achieved by patients with normal renal function. Fampyra must not be administered to patients with mild, moderate and severe renal impairment (see section 4.3).

5.3 Preclinical safety data

Fampridine was studied in oral repeat dose toxicity studies in several animal species.

Adverse responses to orally administered fampridine were rapid in onset, most often occurring within the first 2 hours post-dose. Clinical signs evident after large single doses or repeated lower doses were similar in all species studied and included tremors, convulsions, ataxia, dyspnoea, dilated pupils, prostration, abnormal vocalization, increased respiration, and excess salivation. Gait abnormalities and hyper-excitability were also observed. These clinical signs were not unexpected and represent exaggerated pharmacology of fampridine. In addition, single cases of fatal urinary tract obstructions were observed in rats. The clinical relevance of these findings remains to be elucidated, but a causal relationship with fampridine treatment cannot be excluded.

In reproduction toxicity studies in rats and rabbits, decreased weight and viability of foetuses and offspring were observed at maternally toxic doses. However, no increased risk for malformations or adverse effects on fertility was noted.

In a battery of *in vitro* and *in vivo* studies fampridine did not show any potential to be mutagenic, clastogenic or carcinogenic.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core:

Hypromellose
Microcrystalline cellulose
Colloidal silicon dioxide, anhydrous
Magnesium stearate

Film-coat:

Hypromellose

Titanium dioxide (E-171)
Polyethylene glycol 400

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

The expiry date of the product is indicated on the packaging materials

6.4 Special precautions for storage

Store below 25°C. Store the tablets in the original packaging in order to protect from light and moisture.

After first opening a bottle, use within 7 days

6.5 Nature and contents of container

Fampyra is supplied in either bottles or blister packs.

Bottles

HDPE (high-density polyethylene) bottle with polypropylene caps, each bottle contains 14 tablets and a silica gel desiccant.

Pack size of 56 (4 bottles of 14) tablets.

Blister packs

Foil blisters (aluminium / aluminium), each blister tray contains 14 tablets.

Pack size of 28 (2 blisters of 14) tablets.

Pack size of 56 (4 blisters of 14) tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements.

7. MANUFACTURER

Biogen Idec Ltd
Innovation House,
70 Norden Rd, Maidenhead,
Berkshire, SL6 4AY,
UK

IMPORTER AND LICENSE HOLDER

Medison Pharma Ltd.

10 Hashiloach st, POB 7090, Petach Tikva

Israel

Registration number: 147-80-33543-00

Revised in October 2020

FAMPYRA SPC 0920 V1

