

Patient Leaflet According to the Pharmacists' Regulations
(Preparations) - 1986

This medicine requires a doctor's prescription
Please read the entire leaflet carefully before using this
medicine

DIBENYLIN

Capsules

Composition:

Active ingredient

Each capsule contains:

Phenoxybenzamine Hydrochloride 10 mg

Inactive ingredients

Lactose, talc, titanium dioxide, indigotine,
erythrosine, edible grey ink, gelatin.

Each capsule contains about 180 mg of lactose.

Therapeutic group:

Peripheral vasodilator.

Therapeutic activity:

Treatment of the prostate gland, Treatment of
pheochromocytoma.

When should this medicine not be used?

Do not use this medicine if you are hypersensitive to any
of its ingredients, or to any other medicine that is used to
lower blood pressure.

Do not use this medicine if you had a stroke.

Do not use this medicine if you had a heart attack
(coronary thrombosis) within the last month.

Do not take this medicine without consulting a doctor before starting treatment, in the following cases:

If you are pregnant, planning to get pregnant or
breastfeeding.

If you suffer or have suffered in the past from impaired
function of the: heart and/or vascular system, kidney/urinary
tract, atherosclerosis (including in the brain), or if you suffer
or have suffered in the past from extremity edema. If you
are being treated for shock or your blood pressure is lower
than normal.

How might this medicine affect your daily life?

Use of this drug may cause you to feel dizzy and
therefore caution should be exercised when driving a
vehicle, operating dangerous machinery, or performing
any other activity that requires special vigilance.

Warnings:

If you are sensitive to any type of food or medicine, inform
your doctor before starting treatment with this medication.

This medicine contains lactose. If you are sensitive to
lactose, inform your doctor before taking this medication.

During treatment with this medicine you should be under
medical observation by your doctor, furthermore it is
recommended to take blood pressure readings.

Make sure to get up slowly from sitting/lying position. You
should refrain from ingesting heavy meals and also avoid
increased physical effort with the aim of preventing the
appearance of orthostatic hypotension, dizziness and
palpitations. On the appearance of any of these symptoms,
consult the doctor.

Avoid alcoholic beverages or exposing yourself to high
temperatures.

Drug interactions:

If you are taking any other medications including non-
prescription drugs and nutrition supplements, or if you
have recently finished treatment with another drug, please
notify your doctor in order to prevent any potential risks
or a potential reduction in efficacy that may result from
drug interactions. In particular, medicines of the following
therapeutic groups may interact with your medicine: cough
and cold medicines, medicines that treat heart diseases
and/or blood pressure such as:

- Beta blockers (e.g. atenolol, bisoprolol).
- ACE inhibitors (e.g. captopril, enalapril).
- Calcium channel blockers (e.g. amlodipine, lacidipine).

Side effects:

In addition to the desired effect of the medicine, side effects
may also appear during the course of taking this medicine.
These may include: nasal congestion, fatigue, abdominal

pain or diarrhea.

If these side effects persist and/or are bothersome, please consult your doctor.

The following side effects require special attention:

Blurring of vision, dizziness, fast heartbeat (especially when rising), feeling faint, sexual dysfunction: stop taking the medicine and refer to your doctor immediately!

If at any time you experience side effects that are not mentioned in this leaflet or if you feel an adverse change in your general health, consult your doctor immediately!

Dosage:

Administer according to the doctor's instructions only. Do not exceed the recommended dose.

This medicine is usually not intended for infants and children.

If there is no improvement in your condition within 2-3 weeks refer to your doctor.

This medicine should be administered at set intervals according to your doctor's instructions.

If you forget to take your medicine at the proper time, take the missed dose as soon as you remember, but never take two doses at the same time!

Directions for use:

Do not chew! Swallow the capsule whole with a glass of water, with or after the meal.

How can you contribute to the success of the treatment?

Complete the entire course of treatment as instructed by your doctor.

Even if your condition improves do not stop taking this drug without consulting your doctor.

Avoid poisoning!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning.

If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a doctor!

This medicine was prescribed for the treatment of your ailment; in another patient it may cause harm. Do not give this medicine to your relatives, neighbors or acquaintances.

Do not use medicines in the dark! Carefully check the label and dose each time you take the medicine.

Wear your glasses if you need them.

Storage:

This medicine should be stored below 25°C and in its original package in order to protect from light and moisture. Even if stored properly according to the manufacturer's recommendations, all medicines have a limited shelf life. Please pay attention to the expiry date of your medicine! If you have any doubts, please consult the dispensing pharmacist.

Do not store different medicines in the same container.

Drug registration number: 1260121094

Manufacturer: Dales Pharmaceuticals Inc., UK

Registration holder: Rafa Laboratories Ltd., P.O.Box 405, Jerusalem 91003

The format of this leaflet was determined by the Ministry of Health that checked and approved its content in February 2012.

Dibenylin PIL PB0612-02

794021

021007-G

100302/LF/1

