

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS'
REGULATIONS (PREPARATIONS) - 1986**

Read this package insert carefully in its entirety before using this medicine
This format of this leaflet was determined by the ministry of Health and its
content was checked and approved it in 12/2010

HEMO

Suppositories

Composition

Each suppository contains:

| | |
|--------------------|--------|
| Bismuth Subgallate | 180 mg |
| Benzocaine | 15 mg |
| Zinc Oxide | 120 mg |

Inactive Ingredients: Olive oil, Witepsol w 35 (hard fat), Propyl paraben, Castor oil, Soya bean oil, B.H.T, Sodium methylparaben, Purified water.

Therapeutic activity:

This medicine is intended for the relief of hemorrhoids.

When should the preparation not be used?

Do not use this medicine if you are sensitive to any of its ingredients.

Warnings: Do not use this medicine frequently, or for a long period, without consulting a doctor or if you are pregnant or breastfeeding. If you are sensitive to any type of food or medicine, especially local anesthetics, inform your doctor before commencing treatment with this medicine. Do not use this medicine for 3 days prior to having pancreatic function test.

Drug Interactions: If you are taking another drug concomitantly or if you have just finished treatment with another medicine, inform the attending doctor, in order to prevent hazards or lack of efficacy arising from drug interactions.

Side Effects:

In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine. Treatment with Hemo suppositories may cause change in the colour of the faeces. Such change should not be a cause for worry. In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

Recommended dosage unless otherwise prescribed by your doctor:

Use morning and evening, and after each bowel movement.
Do not exceed the recommended dosage.

This medicine is not intended for administration to children.

Attention:

Do not swallow! This medicine is intended for external use only.

Directions for Use:

Remove the foil wrapper from the suppository. The suppository may be moistened with a some water to aid in its insertion. Lie down on your side and push the suppository deep into the rectum with your finger. Wash your hands thoroughly. Note: If the suppository is too soft, it may be chilled in a refrigerator for about 30 minutes or by placing it under a stream of cold water before removing the foil wrapper. If there is no improvement in your condition within few days, or your condition gets worse, notify your doctor.

How can you contribute to the success of the treatment?

Measures recommended with the use of this medicine:

Proper digestion, prevention of constipation and/or effort on defecation, no lifting of heavy objects.

Avoid Poisoning! This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning. If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a doctor!
Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

Storage: Store below 25°C and a dry place.

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you.

Do not store different medications in the same package.

Licence Number: 028662193200

Manufacturer: Rekah Pharm. Prod. Ltd., 30 Hamelacha St. Holon, Israel.