

PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS REGULATIONS (PREPARATIONS) - 1986
This medicine can be sold without doctor's prescription

CLEAROCIN LEMON, Lozenges **CLEAROCIN CHERRY, Lozenges**

Active ingredients and their quantity:

Each lozenge contains:

| | |
|---------------------|---------|
| Lidocaine | 1.00 mg |
| Tyrosalicylic | 4.00 mg |
| Cetrimonium Bromide | 2.00 mg |

For list of excipients, please see section 6.

Read this entire leaflet carefully before you start taking this medicine. This leaflet contains concise information about the medicine. If you have any further questions, ask your doctor or pharmacist.

This medicine is sold without doctor's prescription. When sold outside the pharmacy (GSL) – the medicine is intended for use in adults and children over 12 years of age. Use in the correct manner. Consult your pharmacist if you need additional information. Refer to your doctor if the symptoms of your illness get worse or do not improve within 5-7 days.

1. What is the medicine used for?

Clearocin is a lozenge for the relief of severe sore throat and mouth infections.

Therapeutic group:

Clearocin contains Lidocaine (local anaesthetic), Tyrosalicylic (antibiotic) and Cetrimonium Bromide (antiseptic).

2. Before you take the medicine:

Do not use the medicine if:

- You are hypersensitive (allergic) to the active ingredients (Lidocaine, Tyrosalicylic and Cetrimonium Bromide) or to any of the other ingredients of this medicine.

Special warnings about the use of this medicine
Before taking Clearocin tell your doctor if:

- You have been suffering from a sore throat for a few days, which is now accompanied by fever
- You have a cut or lesion in the mouth or throat
- Your symptoms do not improve or continue beyond 5-7 days
- The sore throat is accompanied by high fever, dizziness, vomiting or difficulty swallowing

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including non-prescription drugs and nutrition supplements.

Do not brush your teeth immediately before or after using this medicine, as this may reduce the efficacy of this medicine.

Use of this medicine with food:

Do not use Clearocin immediately before or while eating or drinking, as the local anaesthesia may harm swallowing or cause biting of the tongue. Avoid eating while the numb sensation persists.

Pregnancy and breastfeeding:

If you are pregnant, breastfeeding or planning to become pregnant, consult your doctor before using medicines.

3. How to use this medicine

- Check with your doctor or pharmacist if you are not sure.

Usual recommended dosage:

Children over 12 years of age and adults:

For mild - moderate sore throat: suck one lozenge every 2-3 hours.

For severe sore throat: you may suck one lozenge every 1-2 hours.

Do not exceed a dosage of 12 lozenges in 24 hours.

When sold outside of pharmacies, the medicine is intended for adults and children over 12 years of age. This medicine is intended for use for 5-7 days. Permitted for use in diabetic patients.

Do not exceed the recommended dose.

Do not chew or swallow the lozenge.

If you have accidentally taken a higher dosage:

If you have taken an overdose or if a child has accidentally swallowed the medicine, proceed immediately to a doctor and bring the package of the medicine with you.

Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

If you have any further questions regarding the use of this medicine, consult your doctor or pharmacist.

4. Side effects:

Like all medicines, Clearocin can cause side effects, although not everybody gets them. Do not be alarmed while reading the list of side effects, you may not suffer from any of them.

Stop the use of this medicine and refer to your doctor immediately if you suffer from one of the following effects, which can indicate an allergic reaction:

- Difficulty breathing or swallowing
- Swelling of the face, lips, tongue or throat
- Severe skin itching accompanied by a red rash or skin bumps

Additional side effects:

Appear uncommonly (frequency of up to 1:1000):

A tingling sensation in the mouth or throat (discomfort in the mouth)

Appear very rarely (frequency of 1:10,000):

Nausea

If any of the side effects worsens, or if you experience side effects not mentioned in this leaflet, consult your doctor.

5. How to store the medicine

- Avoid poisoning! This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning. Do not induce vomiting unless explicitly instructed to do so by your doctor!
- Do not use the medicine after the expiry date (exp. date) stated on the package. The expiry date refers to the last day of that month.
- Store at a temperature below 25°C and protect from light and moisture.

6. Additional information

In addition to the active ingredients, this product also contains:

Clearocin lozenges – Lemon flavored:

Sorbitol, Citric Acid anhydrous, Talc, Lemon Flavoring powdered, Guar Gum, Quinoline Yellow Lake, Magnesium Stearate, Silicic Acid Colloidal, Saccharin Sodium, Green ZLT 601 Lake, Peppermint Flavoring powdered.

Clearocin lozenges – Cherry flavored:

Sorbitol, Guar Gum, Citric Acid anhydrous, Talc, Polyvinylpyrrolidone, Saccharin Sodium, Magnesium Stearate, Cherry Flavoring, Iron Oxide Pigment Red.

What the medicine looks like and contents of the pack:

Clearocin Lemon are rectangular, dotted, bevel-edged, dark yellow to lemon yellow lozenges
Clearocin Cherry are rectangular, dotted, bevel-edged, pink - red lozenges

Registration holder:

Devries & Co. Ltd., 32 Habarzel st., Tel-Aviv

Manufacturer:

STADA Arzneimittel AG, Stadastrasse 2-18, 61118 Bad-Vilbel, Germany

Drug registration number at the national medicines registry of the Ministry of Health:

Clearocin Lemon: 1382731730

Clearocin Cherry: 1382731729

Revised in May 2021.

DVCLLEPL1120

