

The format of this leaflet has been determined by the Ministry of Health, and its content has been checked and approved in 1.2016

Drontal Plus Flavoured Veterinary

Tablets

For veterinary use only

Must be prescribed by a veterinarian

Composition:

Each tablet contains:

Active ingredients:

Praziquantel 50 mg

Pyrantel Embonate 144 mg

Febantel 150 mg

Inactive ingredients: Please see details in the "Inactive ingredients" section below in this leaflet

Therapeutic group: Anthelmintic

Indications:

Flavoured tablets for the treatment of tapeworms and roundworms in dogs, including the worms types:

Ascarids: *Toxocara canis*, *Toxascaris leonine* (adult and late immature forms)

Hookworms: *Uncinaria stenocephala*, *Ancylostoma caninum* (adults)

Whipworms: *Trichuris vulpis* (adults)

Tapeworms: *Echinococcus* species, *Taenia* species, *Dipylidium* species (adult and immature forms).

Dosage and administration:

According to the veterinarian's instructions.

One tablet per 10 kg bodyweight.

The tablet can be given directly, or disguised in food. Fasting is not required before or after treatment.

Puppies and small dogs:

3-5 kg bodyweight = ½ tablet

Over 5 kg – 10 kg bodyweight = 1 tablet

Medium dogs:

Over 10 kg – 15 kg bodyweight = 1 ½ tablets

Over 15 kg – 20 kg bodyweight = 2 tablets

Over 20 kg – 25 kg bodyweight = 2 ½ tablets

Over 25 kg – 30 kg bodyweight = 3 tablets

Large dogs:

Over 30 kg – 35 kg bodyweight = 3 ½ tablets

Over 35 kg – 40 kg bodyweight = 4 tablets

For the treatment of *Toxocara*, nursing bitches should be dosed two weeks after giving birth, and then every two weeks until the puppies are weaned. Treatment of puppies should be started at age of 2 weeks, and then every two weeks until 12 weeks of age. Thereafter, treatment should be given every three months. It is recommended to treat the puppies and their mother at the same time. Adult dogs should be treated every three months. In cases of severe roundworm infection, a repeat dose should be given after two weeks.

Contraindications and warnings:

Not be used in dogs weighing less than 3 kg.

Do not exceed the recommended dose, especially when treating pregnant bitches.

Do not use simultaneously with preparations containing piperazine.

Consult a veterinarian before treating pregnant bitches for roundworm.

To prevent tapeworm reinfection, take care over the appropriate control of intermediate hosts, such as fleas, mice, etc.

Wash hands well after administering the treatment.

Side effects can be reported to Lidorr Chemicals, by email to:

lidorvet@lidorr.com, or to the Ministry of Health by clicking on the link "Adverse Drug Reactions Report" that appears on the home page of the Ministry of Health web site (www.health.gov.il), which leads to an online form for reporting side effects. Alternatively you can use the following link:

[/https://sideeffects.health.gov.il](https://sideeffects.health.gov.il)

Storage:

Store at room temperature, below 25°C, away from food and out of reach of children.

Do not use this preparation after the expiry date.

Inactive ingredients:

Maize starch, Lactose monohydrate, Microcrystalline cellulose, Povidone, Magnesium stearate, Sodium lauryl sulphate, Colloidal anhydrous silica, Artificial beef flavour irradiated.

Any veterinary medical preparation that has not been used, or any unnecessary material obtained from use of a veterinary medical product, should be disposed of as toxic waste and should not be put into the sewage.

Medicine registration no.: 081 95 92122 00

Manufacturer: Bayer Animal Health, D-51368 Leverkusen, Germany.

Importer and registration holder: Lidorr Elements Ltd., HaNeviim St., Morasha Industrial Area, Ramat Hasharon.