

Veterinary medicine package leaflet
This medicine is supplied on veterinary prescription only
For use in animals only

Marbocyl palatable 20 mg tablets veterinary

Each tablet contains: marbofloxacin 20 mg

Marbocyl palatable 80 mg tablets veterinary

Each tablet contains: marbofloxacin 80 mg

Inactive ingredients:

Lactose, povidone, liver powder, yeast powder, anhydrous colloidal silica, croscopolone, hydrogenated castor oil, magnesium stearate.

Therapeutic group: a third-generation fluoroquinolone group antibiotic.

Medical activity: in dogs - to treat skin infections (for example: Skinfold pyoderma, impetigo, folliculitis, furunculosis cellulites), urinary tract infections (associated or unassociated with prostatitis), and respiratory infections caused by bacteria that are sensitive to marbofloxacin.

Dose: 2 mg/1 kg/day, once a day.

Medium-sized dogs: Marbocyl palatable 20 mg tablet, one tablet for every 10 kg body weight.

Large dogs: Marbocyl palatable 80 mg tablets, one tablet for every 40 kg body weight.

Treatment duration:

- For skin and soft tissue infections, treatment duration is at least 5 days and it can be continued for up to 40 days depending on how the disease progresses.
- For urinary tract infections, treatment duration is at least 10 days and it can be continued for up to 28 days depending on how the disease progresses.
- For respiratory infections, treatment duration is at least 7 days and it can be continued for up to 21 days depending on how the disease progresses.

Contraindications: Do not use in dogs under 12 months old (or under 18 months in especially large breeds who have a longer growth period, such as the Great Dane). Do not use for infections caused by anaerobic bacteria, yeasts, and fungi. Use with caution in dogs who have epilepsy as there is a possibility of neurological side effects. Fluoroquinolones are known to react with orally administered cations such as aluminum, calcium, magnesium, and iron. Bioavailability is reduced in these cases. Low urine pH may reduce the efficacy of this medicine.

Do not use in cats; the recommended doses for cats are lower.

Studies in mice and rabbits did not show any effect of the medicine on gestating animals, but the effect of this medicine was not studied in gestating bitches.

Overdose: May cause serious neurological disorders that are treated symptomatically.

Side effects: No serious side effects were observed at the recommended doses.

Side effects such as vomiting, soft feces, change in thirst pattern, or increased activity may occur. These signs disappear spontaneously after treatment and do not require stopping the treatment.

Young dogs may develop cartilage loss in their joints, so use with caution and dose accurately in young dogs.

You can report side effects to the Ministry of Health by following the link 'Reporting Side Effects of Drug Treatment' on the Ministry of Health home page (www.health.gov.il) which links to an online form for reporting side effects. You can also use this link: <https://sideeffects.health.gov.il>

Warning to the person treating the animal: If you know that you are sensitive to fluoroquinolones, avoid contact with this medicine.

Storage:

- Store below 25°C.
- Keep out of children's reach.

Manufacturer: VETOQUINOL 70204 LURE CEDEX FRANCE

Registration holder: Eliezer Linevitz Ltd., Kanot Industrial Area, Adom 6 st., P.O.B 7006, Israel

Registration number for Marbocyl palatable 20 mg tablets veterinary: 083-92-92396

Registration number for Marbocyl palatable 80 mg tablets veterinary: 083-93-92397

Revised in July 2021 according to MoH guidelines