

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH  
THE PHARMACISTS' REGULATIONS (PREPARATIONS) - 1986**

Read this package insert carefully in its entirety  
before using this preparation

The format of this leaflet was determined  
by the Ministry of Health and its content was checked  
and approved by it in December 2002

# DUO-SCABIL

## Cream

### Composition:

Crotamiton 10%  
Sulphur precipitated 8%

### Inactive ingredients:

Sorbitan monostearate, polysorbate 60, cetyl esters wax,  
cetostearyl alcohol, octyldodecanol, benzyl alcohol, purified water.

### Therapeutic activity:

For the treatment of scabies and itchy areas on the skin  
(pruritis).

### When should the preparation not be used?

Do not use this preparation if you are sensitive to any of its  
ingredients or to other skin preparations for topical treatment.  
Do not use this preparation for acutely inflamed or weeping  
skin, or on cracked skin.

**Do not use this preparation without consulting a doctor  
before starting treatment:**

If you are pregnant.

### Warnings:

If you are sensitive to any type of food or medicine, inform your  
doctor before commencing treatment with this preparation.  
Avoid contact of this preparation with the eyes. In case of  
contact, wash eyes immediately with water.

Nursing mothers should not apply the preparation on their  
nipples or on the surrounding area.

### Drug interactions:

If you are taking another drug concomitantly or if you have just  
finished treatment with another medicine, including non-  
prescription medicines and nutritional supplements, inform the  
attending doctor in order to prevent hazards or lack of efficacy  
arising from drug interactions. This is especially important in  
regards to cosmetic products that may dry out the skin.

**Side effects:** In addition to the desired effect of the preparation,  
adverse reactions may occur during the course of using this  
preparation, for example redness or peeling of the skin.

### Side effects that require special attention:

Local irritation, rash or inflammation (that did not exist before  
using this preparation) - (rare): stop treatment and consult your  
doctor. In the event that you experience side effects not  
mentioned in this leaflet, or if there is a change in your general  
health, consult your doctor immediately.

### Reporting side effects:

Side effects can be reported to the Ministry of Health by clicking  
on the link "Report Side Effects of Drug Treatment" found on  
the Ministry of Health homepage ([www.health.gov.il](http://www.health.gov.il)) that directs  
you to the online form for reporting side effects, or by entering the  
link:

<https://forms.gov.il/globaldata/getsequence/getsequence.aspx?formtype=AdversEffectMedic@moh.gov.il>

In addition, you can report to Perrigo via the following address:  
[www.perrigo-pharma.co.il](http://www.perrigo-pharma.co.il)

**Dosage:** Dosage is according to doctor's instructions only.  
Recommended dosage unless otherwise prescribed by your  
doctor: see section "Directions for use".

**Children's dosage:** In children under three years of age, do  
not apply the preparation more than once a day.

Do not exceed the recommended dosage.

**Attention:** Do not swallow! This preparation is intended for  
**external use only.** Avoid contact of this preparation with the  
eyes and mucous membranes (such as the nose and mouth).

### Directions for use:

Before using this preparation, a warm bath should be taken  
and the body dried thoroughly. Squeeze a dose of cream into  
the hands and rub onto the entire body surface, from the chin  
to the toes, including skin folds and wrinkles (but excluding  
face and scalp), until no preparation remains visible and it has  
been absorbed into the body completely. Repeat this application  
once daily (preferably in the evening) for 3-5 days.

Make sure to change bed linen and underwear every day.  
Thoroughly wash the entire body in a warm bath approximately  
48 hours after the last application.

### How can you contribute to the success of the treatment?

Complete the full course of treatment as instructed by the doctor.  
Even if there is an improvement in your health, do not discontinue  
use of this preparation without consulting your doctor.

**Avoid poisoning!** This preparation and all other medicines  
must be stored in a safe place out of the reach of children  
and/or infants, to avoid poisoning. If a child has accidentally  
swallowed the preparation, proceed immediately to a hospital  
emergency room and bring the package of the preparation  
with you. **Do not induce vomiting** unless explicitly instructed to  
do so by a doctor! **Do not take medicines in the dark!** Check  
the label and the dose each time you take your medicine.  
Wear glasses if you need them.

**Storage:** In a cool place, below 25°C. After first opening can  
be used for 6 months. Even if kept in their original container  
and stored as recommended, medicines may be kept for a  
limited period only. Please note the expiry date of the  
preparation! In case of doubt, consult the pharmacist who  
dispensed the medicine to you. Do not store different  
medications in the same package.

**License number:** 2985.24654

**Manufacturer:** Perrigo Israel Pharmaceuticals Ltd., P.O.B. 16,  
Yeruham.