


PACKAGE LEAFLET

The format of this leaflet was determined by the Ministry of Health and its content was checked and approved by it.



Fenflor
Veterinary

Solution for injection for cattle

1. NAME AND ADDRESS OF THE MANUFACTURER

KRKA, d.d., Novo mesto
Šmarješka cesta 6
8501 Novo mesto
Slovenia



2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Fenflor Veterinary,
solution for injection for cattle

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Each ml of a light yellow to yellow, clear liquid contains:

active ingredient:
Florfenicol.....300 mg

Inactive ingredients:
Propylene glycol
Dimethyl sulfoxide
Macrogol 400

4. INDICATION(S)

Diseases caused by florfenicol susceptible bacteria.
Preventive and therapeutic treatment of respiratory tract infections in cattle due to *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*.
The presence of the disease in the herd should be established before preventive treatment.

5. CONTRAINDICATIONS

Do not use in adult bulls intended for breeding purposes.
Do not use in case of hypersensitivity to the active substance or to any of the excipients.
Do not use in case of resistance to the active substance.

6. ADVERSE REACTIONS

A decrease in food consumption and transient softening of the faeces may occur during the treatment period. The treated animals recover quickly and completely upon termination of treatment.

Administration of the product by the intramuscular route may cause swelling at the injection site which may persist for 14 days. Inflammation at the injection site may persist up to 32 days after administration.

Administration of the product by the subcutaneous route may cause swelling and inflammation at the injection site which may persist at least for 41 days.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.



7. TARGET SPECIES

Cattle.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For treatment:
IM route: 20 mg/kg bodyweight (1 ml/15 kg) to be administered twice 48 hours apart.
SC route: 40 mg/kg bodyweight (2 ml/15 kg) to be administered once only.
Intramuscular and subcutaneous injection.

For prevention:
SC route: 40 mg/kg bodyweight (2 ml/15 kg) to be administered once only.
Subcutaneous injection.

9. ADVICE ON CORRECT ADMINISTRATION

The injection should only be given in the neck.
Swab septum before removing each dose. Use a dry sterile 16 gauge needle and syringe.
The dose volume given at any one injection site should not exceed 10 ml.
To ensure a correct dosage, body weight should be determined as accurately as possible to avoid under dosing.

10. WITHDRAWAL PERIOD

Meat and offal:	by IM (at 20 mg/kg bodyweight, twice):	30 days
	by SC (at 40 mg/kg bodyweight, once):	44 days

Milk: Not permitted for use in lactating animals producing milk for human consumption.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.

Store below 25°C

Shelf-life after first opening the immediate container: 28 days.

Do not use after the expiry date stated on the label.

When the container is breached (opened) for the first time, using the in-use shelf-life which is specified on this package insert, the date on which any product remaining in the container should be discarded should be worked out. This discard date should be written in the space provided on the label and carton.

12. SPECIAL WARNING(S)

For animal treatment only.
Do not exceed the recommended dose.
Do not use in case of hypersensitivity to the active substance or to any of the excipients.
Only administer by the routes outlined under point 8 and 9.
The product should be used in conjunction with susceptibility testing. Official and local antimicrobial policies should be taken into account.

The effect of florfenicol on bovine reproductive performance and pregnancy has not been assessed so it should only be used according to the benefit/risk assessment by your vet.
This veterinary medicinal product should not be mixed with other veterinary medicinal products.

User warnings:

Care should be taken to avoid accidental self-injection.
In case of accidental self injection, seek medical advice and show the label to the doctor.
Do not use the product in known cases of sensitivity to propylene glycol.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

January 2015

15. OTHER INFORMATION

Pack sizes:
50 ml
100 ml
250 ml

Not all pack sizes may be marketed.

Manufacturer: KRKA, d.d, Novo mesto, Šmarješka cesta 6, 8501 novo mesto, Slovenia
License holder: Abic Veterinary Products, P.O.B. 489, industrial Zone, Beit Shemesh
License number: 153-33-33968-00