Patient package insert according to Pharmacists' Regulations (Preparations), 1986 medicine is to be supplied upon obveician's prescription on TIMONII ® 300 RETARD TIMONII ® 600 PETAPD

Prolonged release tablets

Active Ingredient: Carbamazepine 300 mg

For the full list of ingredients see: "Additional information".

Read this entire leaflet carefully before you start using this medicine. This leaflet contains summary information about the medicine. If you have any further questions, refer to your physician or pharmacist. This medicine has been prescribed for the treatment of your linees, Do not pass it on to others. It may ham

This medicine has been prescribed for the treatment of your liness. Do not pass it on to others. It may harm them, even if you think that their liness is similar. This medicine is not intended for children under the age of 6. There are other methods of administration (e.g. swup or immediate release tables) that are more suitable for children under the age of 6.

What is this medicine intended for? Timonii is indicated or the treatment of epipexy specific pain conditions (Trigaminal neuralgia- sharp repetitive facial pain), for treatment of manic disorder and prevention of manic depression- bipolar mood

Therapeutic group:

Before using this medicine

You suffer or have suffered in the past from bore marrow suppression or damage.
You suffer from beat function problems (disvolutificial brock or pontymic).
You are being translet with medicines from the monastimic ordera is holdinor group (MAC) inhibitors) for transmitterior or the superstanding or the suffer sufficient or the superstanding or the sufficient or the s

Special warnings regarding the use of this medicine Since Timonil may cause 'absence seizures' (douded consciousness) or enhance existing 'absence seizures'. It erecommended not to trest batients who suffer from these types of seizures with Timonil.

Before treatment with Timoni I all gour physician 8. Nova allot firms characterized the second seco

You sumer from a socium metabolism disorder, if you nave kidney problems that are related to low socium levels in the blood or if you have kidney problems and you are also taking certain medicines that lower the socium level in the blood (diuretics such as hydrochlorothiazide, furosemido). You suffer or have suffered in the past from cardiac, hepatic or renal function problems (see section 'Side

effects and dosage). - Voi sulfer from degenerative muscular disease (myotonic dystrophy), since problems in cardiac often occur frequently in patients who sulfer from this disease. - Voi have elevised intraccuter pressure (glaucoma) or if you are unable to retain urine.

Additional warnings: Gibler immediately by our physician in order to have a blood coart performed if one of the following signa spepare during featurent with Timenet Sever, sone throat, allergic reactions such as akin relax winn benefing of the hypothesis and the Timese systems, moult use in undersory for hermotians, small medicine, stop the treatment with Timenel Latexy. A start of any moult be and reactions, stop the treatment with Timenel Latexy with Timesel in case of changes in blood coard, maning a treatment of the treatment with Timenel Latexy.

exections, stop be streatment with Thread Hammediathy. However, stop be streatment with Thread Hammediathy. However, stop be streatment with the stream of the stop of the stream of the stop of the stop of the stream of the stop of t

carbamazepine. If you are taking hormonal medicines for treatment of the thyroid gland, your physician will check thyroid function.

o seizurer

to seizures. To prevent undesirable drug interactions (see "Drug interactions") when, in exceptional cases, Timonii The second seco

not given simultaneously. • Due to the possibility of increased skin photosensitivity, protect the skin from strong sunlight during transport with Tamanat

treatment with Timonii. Based on Industrial Considerations, do not change the neticinal between for patients have based on Industrial Considerations, do not change the neticinal between for patients have release or intoxication due to the narrow threspect regime of the active ingredent. Therefore with Timonia chan also a base of activity of the contractive patient and the second second second second second second second second second and the second second second second second second second second a narrow terminal terminal based with indepleted medicine such as carbonargene have had a narrow terminal terminal terminal second second second second second second second activity also of the second terminal terminal second second second second second second second second terminal terminal second second second second second second second second terminal terminal second second second second second second second second terminal terminal second second second second second second second second second terminal terminal second second second second second second second second second terminal terminal second terminal terminal second second second second second second second second terminal terminal second second second second second second second second second terminal terminal second secon

to your obysicial to your physician. • If you are about to undergo any kind of surgery including dental treatment or emergency treatment or any procedure that involves anesthesia, report to the physician that you are taking Timonil.

Children This medicates is usually not intended for children and infants under 6 years of age. There are other methods of administration (such as syrup or immediate release tablets) that are more suitable for children under 6 years of age.

Elderly A lower dosage must be adjusted for the elderly.

Drug interactions Tell your physician or pharmacist if you are taking or have recently taken any other medicines, including non-prescription drugs and nutrition supplements. The dosage may need to changed or other one of the medicines may need to be discontinued. Notify your physician or pharmacit, especially if you are taking.

taking: ine oxidase inhibitors (for depression): Do not take Timonil concomitantly and at least two weeks

Please note that the following reactions may be valid also if the medicine was taken shortly before taking

The effect of Timonil on the plasma concentration of other medicines: Timonil may increase the activity of certain liver enzymes and thereby reduce the levels of other medicines

in the plasma. As a result, the effect of medicines taken concomitantly with Timonil and metabolized in the same way, may be reduced or eliminated. The dosage should be adjusted when using Timonil concentrainty with the foldwing address impedired in affreement methods of administrator: • Antilepteck medicines or ones that are used to treat seizures such as: clonazopam, ethouxivitide, following, privatione, inantidine, inantidine, inantidine, privational ethouses and historials, privational, privational, privational ethouses and historials, privational, privational, privational ethouses and historials, privational ethouse and historials, privational ethouses and Antianxiety medicines such as: alprazolam, clobazam, midazolam.
Medicines for treatment of mental and emotional disorders such a y medicines such as: alprazolam, clobazam, midazolam. i for treatment of mentia and emotional disorders such as: haloperidol, benperidol, clozapine, e, paliperidone, risperidone, queligapine, aripiprazole, zisprasidone. sants such as: buoropion, citalopram, miarserin, mitrazpoine. nefazodone. sertraline. upropion, citaiopram, miarsenn tyline nortriptyline clomipramine Intractions, antitrophyse, nortrighen, contraphyse, comparente. Tetracychie entitholice such as: doscycling) Medicines used to treat (pyshemic) (ingal intections such as: caspolargin, acole type antillangula (such medicines used to treat (pyshemic) (ingal intections such as: caspolargin, acole type antillangula (such medicines for treatment of vial diseases or ADS such as Indinny', thoran's, sagutaner, Medicines for treatment of vial diseases or ADS such as Indinny', thoran's, sagutaner, Medicines for treatment of alternities, isolandiscut, Medicines for treatment of alternities, isolandiscut, Medicines for treatment of alternities, indinnetscut, Medicines for treatment of antiener of such as thorange, instructions, Medicines for treatment of cardies antiphysics, such as cardiner, Medicines for treatment of cardies antiphysics as charged inter, Medicines for treatment of cardies antiphysics and as cardiner, Medicines for treatment of cardies antiphysics and as cardiner, Medicines for treatment of cardies antiphysics and as cardiner, Medicines for treatment of cardies antiphysics and as cardiner, Medicines for treatment of cardies antiphysics and antipacting. trazodone, imipramine, amit edicines for treatment of heart disease: digoxin, ivabradine sta blockers for lowering blood pressure such as: proprano

Belt biodens for lowering blood pressure such as propresed. Medicines for weight biod pressure such as information and the second pressure and the second pressure and the second pressure such as information, weight the second pressure such as infrarediate, proceeding and the second pressure such as infrarediated pressure such as infrarediated pressure and the second pressure is (normonal contraceptives). Medicine for treatment of impotence: tadalafi Medicine for treatment of breast cancer: toremifene

Taking contraceptive pills may lead to a loss of contraceptive effect and thus to unwanted pregnancy, as well as sudden intermenstrual bleeding. Therefore, consult your physician about using alternative methods

The plasma concentration of phenytoin can be either reduced or increased by carbamazepine, which can lead in exceptional cases to states of confusion or even come.

Timonil may lower plasma levels of the medicine bupropion (a medicine used to help stop smoking) and increase its degradation rate, thus reducing clinical efficacy and safety of bupropion.

Timonil may lower plasma levels of the medicine trazodone (for treatment of depression). However, it appears that Timonil enhance the depression tranquillising effect.

Timonil may accelerate the degradation of zotepine (treatment of mental health problems)

Medicines that decrease the plasma concentration of carbanazepine • Other antiepilepic medicines such as: ethosuminds, oxcarbazepine, phenobarbital, phenytoin, prindrone, programing, particular and possible (conceptant) • Medicines for treatment of eavere respiratory diseases such as: theophylline, antinophylline. • Antibiotic: reframential of earer. bioethroin.

otic: rifamploin. ines for treatment of cancer: cisplatin, doxorubicin. bia monificiane used to treat decression: Hypericum perforatum- St. John's Wort

Alternatively, the plasma levels of the active metabolite of TinonII may be increased by valproic acid and

comitant administration of felbamate may reduce the carbamazepine levels in the plasma and the level of the carbamazepine metabolite; at the same time the levels of felbamate may be

Due to these interactions, it is recommended to monitor the plasma levels, especially during concomitant treatment with other antisplieptics, and to adjust the dosage of Timonil if necessary.

A decrease in the carbamazepine plasma concentration may cause an aggravation of the disease symptoms, as well as cause recurrence of epileptic seizures, seizures in multiple scienceis or recurrence of facial pain or pain in the corpharyngeal space.

Biold plant or part in the do-purpress seture. Medicines tay Loncense to plasma concentration of carbamazepine: Medicines tay Loncense to plasma concentration of carbamazepine: • extendences for treatment of bacterial infections are at coprolocatin, manotile antibiotics such as: • professions for treatment of functional infections or antibiotics of the acole group, such as • Medicines for treatment of functions control treatment of the acole group, such as • Medicines for treatment of the acole seture of the acole group, such as • Medicines for treatment of the acole seture of the acole provide the acole seture of the acole • Medicines for treatment of the acole seture of the acole provide the acole seture of the acole of the acole seture of the acole of the acole aco

ants such as: one

te relaxants such as: oxybutynin, danuolene.

Vedicines for treatment or timesus environment. Anticeaquiants such as tickopidine. Analgesics such as: Ibuprofen, destropropoxyphene/propoxyphene. Analgesics such as: Ibuprofen, destropropoxyphene/propoxyphene. Analgesics such as: Ibuprofen, destropropoxyphene/propoxyphene.

flavoxamine. Medicine to inhibit secretion of sexual hormones (gonadotopins): danazol. Medicine for treatment of gastrointestinal ulcers: cinnetidine. Medicine for treatment of neathrum: cinneprazio. High dosage nicotinamide (vitamin B). Medicines for treatment of allergic reactions such as: terferadine, loratadine.

An increase in the plasma concentration of carbarnazepine may enhance the side effects (dizziness, fatigue, walking unsteadily, double vision). If these symptoms occur, the plasma concentration of carbarnazepine must be checked and the doscene reduced if necessarv.

Medicines that increase the plasma concentration of the active metabolite of carbamazepine restaring una inclusion de passia concernación de la aure menaciate ol carbanazario carbanazarione 10.11-egocióde: Other antiepileptic medicines such as: oxcarbazepine, primidone, progabide, valproate, valnoctamide. Medicines for treatment of mental and emotional disorders quetaaine. Ioxabine.

Additional medicines:

dottoma medicines: Concomitant use of carbamazepine and levetirscetam may increase carbamazepine toxicity. The concomitant use of Timonil and lithum (a medicine used for treatment of depression an bipolar disorder), neuroleptics: medicines used for treatment of mental and emotional disorders) metocloppranide (a medicine used for treatment of particular disorders) may lead to neurologic ride officier rido effecte

In the case of patients who are treated with neuroleptics it is important to remember that carba In the case of paramaterial who are reasonable in the original in an approximate or internet house the acceleration of the energical cases of the second cases of the Note in particular that concomitant use of lithum (a melicine used for treatment and prevention o certain mential and psychological discortes) and Timoni may enhance the damaging effect of these two active ingredients on the nervous system. It is therefore necessary to monitor plasma levels of the two melicines. Treatment with neurophycitics should be discontinued at least 5 weeks before treatment with Timonii, and not be given concomitantly. Pay attention if the following symptoms appear: walking unstability discussion, convisions or threating anys (introtant inystagmus), a henhand reflexes, musci na levels of the

beliching. In the literature there are indications that taking Timontil in additional to antipsycholics may increase the risk of neurolegic malignant syndrome (lite-threatening coddition with increase in body temperature and muck stiffless). Solven-Johnson syndrome (lite threatening akk neation). The hepatoboxicity of isonizad (medicine for treatment of taberculosis) may be increased by Timonil. The concentration us of Timonil and some diverties such as hydrochrochrothacia and furosemide man

Ingermal neurange: Maintaining the maintenance dosage suitable for treatment of pain for a period of a few weeks has been proven effective. It is recommended to carefully reduce the dosage to determine if the pain has passed during the treatment period. If the pain persists, continue the treatment with the maintenance dosage. Prevention of certain psychiatric disorders (preventing manic depression): e a decrease in blood sodium

cause a decrease in blood sodium levels. The effectiveness of muscle relaxants such as: pancuronium may be damaged by concomitant use of **Timonil**. For this reason, these medicines may clear faster from the body. Carefully monitor patients The eff

treated with muscle relaxants and increase the dosage if necessary. • When taking Timenil concomitantly with isotration (medicine for treatment of acne) plasma levels of carbanazapien must be monitored. • The concomitant use of Timenil and paracetanol (analgesic and antipyretic) may reduce the bioavailability and effectiveness of paracetanol. Long-turne use of Timenil and paracetanon may lead Tests and follow up: Based on the side effects lated in section "side effects" and the hypersensibility reactions, it is necessary to perform a routing blood count during heatment with "Temorit to negularly check here and kidney functions, perform a routing of the section of the count of the section of the Carbon of the section o age. Sine appears to enhance the elimination of hormonal medicines for treatment of the thyroid

Please refer to your physician or pharmacist if you are under the impression that the effect of TimonII is

If you forgot to take this medicine at the specified time, do not take a double dose. Take the next dose at the regular time and consult a physician. Make sure to take the medicine at the right times to

Even if there is an improvement in your medical condition, do not stop taking the medicine without consulting a physician. Suddenly stopping to take the medicine may harm the success of the treatment and cause recourrence of the excitences.

Do not take medicines in the dark! Check the label and the dose each time you take a medicine. Wea

If you have any further questions regarding the use of this medicine, consult your physician or pharmacist.

Side effects Like any medicine, the use of **Timonil** may cause side effects in some users. Do not be alarmed while reading the list of side effects. You may not suffer from any of them.

Refer Immediately to a physician if one of the following side effects appears. These may be an early sign for severe damage in the circulatory system, liver, kidneys or other organs and may require medical

sign for somer damage in the circulatory vysion, live: Klenny or other drags and may require models attendent immediately memory and the rate score house, rate, model visions, reselling in the hyroth-nodes. Increases its association is incluing (maintende by changes in blood court, especially by co-decreases in the municed while blood courts on second that remain, now blood (signs of changes). A sign, including a standard in the standard standard in the standard standard standard blood court, maintender while blood courts and standard blood (signs of changes). A spearsment of system, set and standard in the like, isocially and standard standards. A spearsment of system, set and standard in the like, isocially and the standard in the standard blood court, maintender of system and the standard standard standard standards. A spearsment of system, set and maintain on the like, isocially associated and the standard standard standards. B of standards and standards and standards and standards and standards and standards and standards. B of standards and standards and standards and standards and standards and standards and standards. B of standards and standards and

eaction). Apathy, confusion, muscle twitching or significant worsening of convulsions (symptoms that may be Apaimy, contusion, muscle whiching or significant worsering of convusions (symptoms that main makes to be be board online) which if mode and actives sensitivity to light (large of manning)).
Nuscle attifness, high fores, atterned consciourness, high blood pressure, excessive salivation (sign munorlepit) multipart synchrone and twenters.
Versida the heartbeat and check pain.
Versomet attachances and weakness.

Additional side effects If was observed that the risk of side effects was lower when Timonii was taken as a monotherapy as

It was observed that the risk of side effects was lower when Timonil was taken as a monoherapy as opposed to combined treatment with whore natiopiliped medicines. Some of the side effects are dosage desendent and usually occur at the beginning of treatment, when the initial dosage is to high, or in adults. These side effects include in the contril nervous system disorders (dizziness, headache, walking disorders, drawsiness, sedation, fatgua, double vision, accommodation disorders such as buined vision), gastrichtefand adorders (nausea and vomithg) and allerigt skin

dosage should be gradually increased, if possible. Central nervous system disorders may be a sign of a relative overdose or large fluctuations in the medicine plasma levels. It is therefore recommended to check the plasma concentration of the medicine.

In certain cases, a decrease in bone density (osteoporosis- causing fractures) was reported. Please consult your physician or pharmacist if you are taking antiepleptic medicines for a long time, if you are suffering from osteoporosis or you are taking actistone or other steroid hormones.

Inversion bisoleters. Nausea and vorniling. Severe allergic skin reactions, with or without fever and hives (urticaria). Increase in blood enzyme values that indicates liver function - Gamma-glutamyl transferase (gamma-GT).

Common side effects (affects that occur in 1 - 10 out of 100, users): • Water accumulation in the tissue (overna), recluded fluid excretion, weight gain, hyponatremia (reduc blood sodium level) and docreased plasma comolally, which may ranely lead to water indocade manifested by tadjue, vomilian, hestache, confusion states and other neurological disorders that c

¹ increase in taked incident takes an end of the second in the second in takes a first and second in takes a first and second. The second is taken a second in takes a second second in takes a second s

Heart conduction disturbance (vv biocx), sometimes accompanied by loss or consucuences and main or low blood pressure. Cocasionally to ranely stowing of the heart beat (bradycardia) and arrhythmiss, circulatory system failure, heart failure and worsening of pre-wsking coronary heart disease, phelotis and blood cot formation.

scaly takin inflammation where endering of the whole body and skin pelling appart (expfrredema).", Kollany disorders such as exercised of protein in the unine, blood in the unine (haematuria), decreased unine production (oliguria) or elevated bood urea nitrogen (azotaemia). Increase in blood enzyme values that indicate liver function (manaminases).

Rare side effects influence and receive inter surction (transaminases). * A number of severe and life-finataming skin reactions (Stevens-Johnson syndrome and toxic epidermal necrohis) (see section "Additional warnings"). • Umphademostry.

Inflammation (dermatitis) where the skin or mucous membranes peel off (exfoliative dermatitis).

ness, dizziness, undesirably strong tranquility, fatigue, somnolence, exhaustion, walking and

These side effects disappear on their own within a few days or after a dosage reduction. The

Very common side effects (effects that occur in more than one out of 10 users); • Changes in blood count, such as a decrease in the number of white blood cells (leukopenia); • Drowginges, diprivates understands, decrease in the number of white blood cells (leukopenia);

eau to: ● Headache, double vision, accommedation disorders (such as blurred vision). ● Loss of appetite, dry mouth ● Increase in blood enzyme values that indicate liver function - alkaline phosphatase

dose at the regular time and consult a physician. Make sure t maintain the efficacy and safety of the treatment. Continue with the treatment as recommended by the physician.

ses if you need then

Side effects

reaction).

duced if necessary. mended to perform a blood count and test the liver functions before treatment with Timonii. then

o strong or too we

Folic acid deficiency.
Hallucinations (seeing and hearing things that do not exist), mood changes such as deprively marks table (elevated mood accompanied by aggression), loss of appetite, restlessner

aggressive behaviour. Novement disorders such as involuntary movements of the mouth and face causing face distortion, bolted novements, speech problems (situred speech), paraesthesia, peripheral neuropathy (polyneuropathy),

Abdominal pain. Henetilie or liver inflammation (various forms of henetilis) and liver disease with destruction and Hepatitis or liver inflammation (various forms of hepatitis) and liver disease with destruction and disappearance of the intrahegatic biel ducts. Acute hepatitis or life threatening liver failure particularly in the first months of treatment. Rohing or a chronic inflammatory disease of the connective tissue that damages the skin and is manifested by a not rank with acutes on the face (topular exprementacus).

Mary area side effects (effects that coccur in less than consultances area col of 10000) (grannadocroso), splately ameres, perrofroponia (simultances alerces in red and white bood cells and as patients, patient or maneria, perrofroponia (simultances alercess in red and white bood cells and area patients), splately ameres, perrofroponia (simultances alercess in red and white bood cells and area patients), splately ameres and of cells, and area of the forms of anaemia, reliculorybrais and versional forms of portphylin.

various torvine of portphysis. Acutal general alloys reaction and assiglto maniholitis accompanied by muscle liviticity and profileration of cardin while allowed calls (assignt) analphysical, swelling of kin and muscle allogibadem-consense in portacities and analysis of the male beast or mitige production unrelated to 15th of theast-fielding in women. The state of the state of the state of the state of the male beast or mitige production to the state of the state of the state of the state of the male beast or mitigeness of the state of the the state of the the state of the the state of the of the state of the of the state of the st

nyiunctMMs, cataract and increased intraocular pressure. aring disorders such as timults, exaggerated or decreased auditory sensitivity, change in pitches. persensitivity of the lungs accompanied by fever, breathing difficulties, pneumona (pneumon wellis-chronic inflammation of the alwedi walla).

arevenues-crinonic imammation of the alveoli walls). Inflammation of the oral and pharyngeal mucose (stomatilis, gingivitis, glossitis), pancrestitis. Lipht sensitivity, skin rednese accompanied by different forms of skin rash and subcutaneous nodules, skin rash due to bleeding of small blood vessels (capitilaries) under the skin (purpura), hair loss, increased sweating, charges in skin signmentation, acre, hinsulism (excess male type hair in women), vasoutat

International and a second sec

Junings in the tone training matrixe.
Junings in the tone training matrixes are associated with the tone of the interface.
Decrease in vitamin 10, levels and index to both homosysteine levels.
Decrease in vitamin 10, levels and index to both homosysteine levels.
Section 20, levels and index to both homosysteine levels.
Section 20, levels and index to both homosysteine levels.
Section 20, levels and index to both homosysteine levels.
Section 20, levels and index to both homosysteine levels.
Section 20, levels and levels are associated and levels.
Section 20, levels and levels are associated and levels.
Section 20, levels and levels are associated and levels are associated and levels are associated and levels.

caresmanagement have been regorded. Unbrugement parts causes or two-jubic parts searches when characterization of the search of the Stor reactions and provide the search of the search of the search of the search of the Stor reactions accompanied by rays, then the search of the search of the search search of the Stor reactions of participants in the search of the search of the search of the Part or search of participants when the search of the search of the search of the Part or search of participants when the search of the search of the search of the search of the Part or search of participants when the search of the

Reactivation of the herpes virus interction (human impression of time) to the second of the herpes virus and the second of the second of the days of become a bothersome, refer to your physician.

Proceeding to all relaxations are a decreased another to write above being being includence and the second If you experience a side effect, if one of the side effects worsens, or if you suffer from a side effect not mentioned in the leaflet, consult your physician.

Reporting side effects: Side effects can be reported to the Ministry of Health by clicking on the link "Report Side Effects of Drug Side effects can be found on the home page of the Ministry of Health website (www.health.pou/)) which enforcement that can be found on the home page of the Ministry of Health website (www.health.pou/)) enforcement page side the side of the Ministry of Health website (www.health.pou/)) enforcement page side the side of the Ministry of Health website (www.health.pou/)) enforcement page side of the side of the side of the Ministry of Health website (www.health.pou/)) enforcement page side of the side of the

Avoid poleoning! This medicine, and any other medicine, must be stored in a closed place out of the reach of childen and/or infants, to avoid poleoning. Do not induce vomiting unless explicitly instructed to do so by the physician! The medicine after the explive (soc. date) stated on the package. The expiry date refers

In addition to the active ingredient carbamazepine, the medicine also contains: Methacrylic acid-ethyl acrylate copolymer (1:1), Ammonio methacrylate copolymer (type B), Cellulose, Microcrystalline, Sodium starch glycolate (type A), Taic, Purified water, Silica colicidal anhydrous,

What the medicine looks like and contents of the package:
Timonil 300 mg-Protonged release tablets, white, round and flat, cross-scored on one side and engraved
with a 'T on the other side. Available in bifer packages of 50, 100 and 200 tablets.

Timonil 600 mg- Prolonged release tablets, white, oval, biconvex tablet with a score line on one side and engraved with a 'T' on the other side. Available in bitster packages of 50,100 and 200 tablets

Drug registration number at the national medicines registry of the Ministry of Health: Timonil 300 mg: 068-94-27055

Registration holder and his address: Megapharm Ltd. P.O.Box 519, Hod Hasharon 451050

Manufacturer and his address: Desitin Arzneimittel GmbH, Hamburg, Germany

This leaflet was checked and approved by the Ministry of Health in January 2016.

*1 According to the literature, a decreased amount of white blood cells (benign leukopenia) is very corr

Side effects with unknown frequency (effects whose frequency has not been determined yet): Allergic cross-reactions with other antispipelpic medicines.
Syndrome characterized by sidn reactions, sevial in mych nodes, fever and involvement of other organs such as DRESS (Drug rath with ecsimophilis and systemic symptoms syndrome).
Sikin reaction accomparated by redness and bistering (acute generalized rash - AGEP).
Damage to the bene marrow.

weolitis- chronic infl

rting side effects

How to store this medicine

ure below 25%

to the last day of that mon a at a terr

Additional information

Timonil 600 mg: 067-24-27139

Timonii Pil. P80516-06

lagnesium stearate,

Carbamazapine appears to enhance the elimination of horminolial medicines for treatment of the thytoic gland and therefore it is necessity to increase the docage of these modificies in patients who suffer from end of the treatment with Timonia. The docage of these medicines may need to be adjusted. Concomitant use of Timonii With antidepressants that inhibit servicini absorption such as fluxewine may cause a toxic syndrome of serotonin accumulation cross a wwell during the frst month of heatment and later once a month. After six months of treatment is is enough to perform this test twole to four times a year. Furthermore, it recommended to perform urine tests and measure urine urea nitrogen levels before treatment with carbamazepine and regularly during the treatment with carbamazepine.

cause a toxic synarome of serotonin accumulation. Antidepressant, nefazodone. It is recommended not take this medicine with Timonil. Carbamazepine may cause a decrease in the blood concentration of this medicine and even loss of its effect. Furthermore, the level of the active ingredient carbamazepine will increase and the level of the active metabolite will the heatment will cathemization. If you have accidentially taken a higher desage you may experience the side effects described in sector take effects in a non-server burn. You may experience the blowing prophone. The sector takes and the sector takes and the sector takes and the sector takes and takes and the sector takes and the sector takes and the sector takes and takes and takes and the sector takes and takes takes and takes takes and takes takes and takes takes and takes takes and takes takes and takes takes and takes takes and takes takes and takes takes and takes takes and cecrease. Concomitant use of carbamazepine and other medicines that may cause cardiac conduction disturbance

Concomitant use of carbamazappine and other medicines that may cause cardiac conduction disturbance, aniamtrythmic medicines, certain anidyersessants (cycic) antidepressants) or enythromytoin (antibiotic) increases the risk of cardiac conduction disturbances. Antisejöspic medicine: seliciant-zegoria exostate. Concomitant use with carbamazapine may increase esilicatrazprine acetate toxidy. Double vision, coordination disturbances and dizintenses are direct common during concomitant use of esicatrazpepine acetates and Timotil than during concomitant use with common during concomitant use of esicatrazpepine acetates and Timotil than during concomitant use with common during concomitant use of esicatrazpepine acetates and Timotil than during concomitant use with common during concomitant use of esicatrazpepine acetates and Timotil than during concomitant use with common during concomitant use of esized than the setting concentration disturbances.

ing concomitant use or earch esilicarbazepine acetate and other anticonvulsants. • Oxcarbazepine: may cause a decrease of 0 - 22% in the plasma level of the active ingradient when used with carbamazepine, with a simultaneous increase of 30% in the carbamazepine epoxide level.

Use of this medicine and food

There have been several reports of an increase in the blood level of the active ingredient carbamazepine after drinking grapeful iluice.

Use of this medicine and alcohol consumption Use any other substances that affect the onthal nervous system. Timenil may reduce alcohol tolerance. Avoid alcohol consumption during the treatment with Timenil since the effect of Timenil may change and enhance unpredictably by alcohol.

Pregnancy If you are proposed on the pregnance, planning to become pregnant or breastfeeding, consult your physical proposed on the pregnance of the pregnancy. Carbomacopine may be prescribed during pregnancy only after the physican carefully evaluates the

Cabital science of the formation of the analysis of the program by the source of the program by the source of the

Breastfeeding The active ingredient carbamazepine passes into breast milk in low quantilies. However, the benefits of breastfoeding abuild be weighed against the risk of aide effects in the infant. Increfere, consult your physician before starting to breastfeed. If the infant's weight gain is insufficient, the infant is not sufficiently airt, the develops allering skin reactions or there is liver damage, stop the breastfeeding.

Driving and use of machinery As a result of table effects such as discretes, somorisence, forgun, unstandinerse, double vriton, As a result of table effects such as a discrete effects and the second second

will not be able to react quickly and efficiently to sudden occurrences. Therefore, be careful while driving a vehicle, operating machinery and every activity that requires attention. Do not work in a place where there is no safe place to stand. Especially note that alcohol consumption during treatment increases the impairment in driving ability! As for children, they should be warmed about riding bicycles or playing near roads etc.

use according to your physician's instructions. Check with the physician or pharmacist if you are

The second secon

Administration: The labeles can be divided. The labeles can be divided. It is possible to discove the tablet in a glass of water (as a suspension) and drink the suspension immediately afterwards.

Duration of treatment and discontinuing it: The duration of treatment is determined by the physician and depends on the indication for which the medicine was prescribed and the personal response of the patient.

reatment is fundamentally a long-term treatment. A specialist with experience in Artigologics featiment is inclamantally a long-term featiment. A specialist with experience in Treats Artigologics featiment is inclamantally a long-term featiment. A specialist with experience in Treats foremative, reducing the docage of discontinuing treatment will be done only if the patient has not suffer from secures for at least two or three years. To children – reducing the docage can be done sub-the docage valid the docage valid terminal under the children – reducing the docage can be done in such a way that the docage will terminal under the children – reducing the docage can be done in such a way that the docage will terminal under sub-type docage can be done and the docage terminate terminate terminate activity detected to do detectorist.

abnormal spermatogenesis were repo

Fertility sexual dysfunction, e.g. erectile dysfunction or decreased libido, as well as decreased male

Driving and use of machinery

How to use this medicine

Do not exceed the recommended dose

Trigeminal neuralgia: