

The format of this leaflet was determined by the Ministry of Health and its content was checked and approved in October 2016

## 1. NAME OF THE VETERINARY MEDICINAL PRODUCT

Engemycin Spray Veterinary

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL contains

Active substance:

Oxytetracycline hydrochloride	25.00 mg
(equivalent to oxytetracycline	23.15 mg

Excipients:

Patent blue V (E131)	1.25 mg
----------------------	---------

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Cutaneous spray, suspension

Green to green-blue suspension

## 4. CLINICAL PARTICULARS

### 4.1 Target species

Cattle, sheep and pigs

### 4.2 Indications for use, specifying the target species

For the treatment of the following infections caused by, or associated with, organisms sensitive to oxytetracycline in cattle, sheep and pigs:

- Treatment of foot infections caused in particular by: *Dichelobacter nodosus*, *Fusobacterium necrophorum* and other *Fusobacterium* spp., and *Bacteroides* spp.
- Supporting treatment of superficial wound infections following surgery or physical injuries, e.g., tail biting in pigs, scratches and abrasions.

### 4.3 Contraindications

Do not use for treatment of teats in order to prevent the product from getting into milk.

Do not use in animals in cases of hypersensitivity to oxytetracycline or to any of the excipients.

#### **4.4 Special warnings for each target species**

None.

#### **4.5 Special precautions for use**

##### **i) Special precautions for use in animals**

The animals should be treated in a well ventilated area.  
Do not spray in or near the eyes.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

##### **ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Because of the risk of sensitisation and contact dermatitis, the user should avoid skin contact. Wear appropriate impermeable gloves whilst handling the product.

Because of risk of eye irritation, contact with the eyes should be avoided.

Protect the eyes and face.

Do not spray on a naked flame or any incandescent material.

Do not pierce or burn the container, even after use.

Avoid inhaling vapours.

Apply the product in the open air or in a well ventilated area.

Wash hands after use.

Do not eat or smoke whilst administering the product.

In case of accidental ingestion or in case of contact with eyes, seek medical advice immediately and show the label to the physician.

#### **4.6 Adverse reactions (frequency and seriousness)**

None known.

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Any suspected adverse events should be reported to the Ministry of Health according to the National Regulation by using an online form

<http://forms.gov.il/globaldata/getsequence/getsequence.aspx?formType=AdversEffectMedic@moh.gov.il>

#### **4.7 Use during pregnancy, lactation or lay**

Can be used during pregnancy and lactation.

#### **4.8 Interaction with other medicinal products and other forms of interaction**

None known.

#### **4.9 Amounts to be administered and administration route**

For topical use only.

Shake well before use. The spray container is suitable to be used in upright and inverted positions. Before application properly clean the surface to be treated, spray the product for 1-2 seconds, at a distance of 15-20 cm, until the area has a homogeneous colour. Repeat the treatment every 12 hours for 1 to 3 days, depending on the healing process.

To achieve the best results in case of pedal lesions the following instructions are recommended:

- fully clean the foot area, completely removing foreign material, exudates and necrotic tissue
- keep the animal on dry ground for 12 hours after each application.

#### **4.10 Overdose (symptoms, emergency procedures, antidotes), if necessary**

None known.

#### **4.11 Withdrawal period(s)**

##### **Cattle, sheep:**

Meat and offal: Zero days

Milk: Zero hours

##### **Pigs:**

Meat and offal: Zero days

Stained part of the pig skin must be removed prior to the rest of the animal being used for human consumption.

### **5. PHARMACOLOGICAL PROPERTIES**

Pharmacotherapeutic group: antibiotics for topical use, tetracyclines

ATCvet code: QD06AA03

#### **5.1 Pharmacodynamic properties**

Oxytetracycline is produced by fermentation of *Streptomyces rimosus*.

It possesses broad spectrum antimicrobial activity against a wide range of Gram +ve and Gram -ve bacteria including target pathogens *Dichelobacter nodosus*, *Fusobacterium necrophorum* and other *Fusobacterium* spp., and *Bacteroides* spp.,.

Oxytetracycline is bacteriostatic and acts by inhibiting protein synthesis within the cell.

## **5.2 Pharmacokinetic particulars**

When administered topically, oxytetracycline absorption is negligible and the drug comes into direct contact with bacteria on the skin and in superficial lesions on external body surfaces. The marker dye indicates the extent of the treated area.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Patent Blue V (E131)  
Polysorbate 80  
Isopropyl alcohol  
Mixture of hydrocarbons on butane basis (LPG), with denaturant

### **6.2 Incompatibilities**

None known.

### **6.3 Shelf life**

Shelf-life of the veterinary medicinal product as packaged for sale: 2 years.

### **6.4. Special precautions for storage**

The product should be stored at a temperature below 25°C. Keep away from fire or sources of ignition. No smoking.

### **6.5 Nature and composition of immediate packaging**

Pressurised lacquered aluminium spray container containing in each 200 ml pack 5 g oxytetracycline hydrochloride and a blue colourant. The spraying valve consists of lacquered tinplate and different plastic materials and enables the container to be operated in upright and inverted positions.

### **6.6 Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products**

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

## **7. LICENSE NUMBER**

157-22-34471-00

**8. MANUFACTURER**

Intervet Productions SRL, Via Nettunense Km.20.300, Aprilla (LT) Italy.

**9. MARKETING AUTHORISATION HOLDER**

Intervet Israel Ltd., Neve Ne'Eman Industrial Park, Hod HaSharon 45240