הודעה על החמרה (מידע בטיחות) בעלון לרופא מעודק 05.2013)

תאריך: **08 פברואר 2017**

שם תכשיר באנגלית ומספרי רישומו : <u>טרג'נטה 5 מ"ג טבליות 149-96-33738-00</u>

שם בעל הרישום: <u>בורינגר אינגלהיים ישראל</u> ההחמרות מסומנות על רקע <mark>צהוב.</mark>

ההחמרות המבוקשות		
טקסט חדש	טקסט נוכחי	פרק בעלון
Changes in laboratory findings were similar in patients treated with TRAJENTA 5 mg compared to patients treated with placebo. Increase in Uric Acid: Changes in laboratory values that occurred more frequently in the TRAJENTA group and ≥1% more than in the placebo group were increases in uric acid (1.3% in the placebo group, 2.7% in the TRAJENTA group). Increase in Lipase: In a placebo-controlled clinical trial with TRAJENTA in type 2 diabetes mellitus patients with micro- or macroalbuminuria, a mean increase of 30% in lipase concentrations from baseline to 24 weeks was observed in the TRAJENTA arm compared to a mean decrease of 2% in the placebo arm. Lipase levels above 3 times upper limit of normal were seen in 8.2% compared to 1.7% patients in the TRAJENTA and placebo arms, respectively.	Changes in laboratory findings were similar in patients treated with TRAJENTA 5 mg compared to patients treated with placebo. Changes in laboratory values that occurred more frequently in the TRAJENTA group and ≥1% more than in the placebo group were increases in uric acid (1.3% in the placebo group, 2.7% in the TRAJENTA group).	6.1 Clinical trial expertise